

## **RESPONSE OF GROWTH, PRODUCTIVITY AND STORABILITY OF GARLIC (*Allium sativum* L.) TO FOLIAR SPRAY WITH MAGNESIUM AND YEAST EXTRACT**

**El-Morsy, A. H. A; U. M. Saif El-Deen and A. S. Ezzat**

**Veg. Res. Dep., Hort. Res. Inst., Agric. Res. Center, Giza, Egypt.**

### **ABSTRACT**

Foliar application of mineral nutrients considers a method of supplying nutrients to higher plants more rapidly than methods involving root application, which makes the nutrients more efficient. It is a quick and efficient method of supplying micro elements in particular. It can, also be used to satisfy acute needs of macro nutrients. This study was conducted on garlic cv. Sids-40, in the private farm at Kafr Meet Faris village, near El-Mansoura, Dakahlia Governorate, during 2008/2009 and 2009/2010 seasons to study the effect of foliar application with certain magnesium concentrations (0.0%, 0.2%, 0.4% and 0.6% as Mg-citrate) either single and/or in combination with foliar application of some yeast extract concentrations (0.0, 25 ml/L, 50 ml/L and 100 ml/L) on plant growth, yield and its components, as well as chemical constituents and storability of bulbs during the storage period.

#### **The obtained results could be summarized as follows:**

In general, results showed that the plants sprayed with Mg-concentrations were better than those of the unsprayed ones. Increasing the foliar applied magnesium concentration from 0.2% to 0.6% Mg significantly increased plant height, number of leaves/plant, plant dry weight and bulbing ratio as well as total yield and bulb weight and diameter. Moreover, foliar application of magnesium at 0.6% Mg significantly increased concentrations of N, P and K in cloves. In addition, TSS% and volatile oils were increased. This concentration had the most interesting observation in the enhancing of storability. On the other hand, foliar application of yeast extract at 50 ml/L gave rise to significant increases in plant height, number of leaves/plant; plant dry weight, bulbing ratio, total yield, bulb weight and diameter and clove weight as well as chemical constituents in cloves and decreasing bulb weight loss percentage during the storage period compared with the other treatments. The combined treatments of Mg-concentrations and yeast extract were generally more effective on the most studied parameters than single ones. The best results were obtained by foliar application of 0.6% Mg with foliar application of yeast extract at 50 ml/L. This treatment achieved increases in yield at the end of the storage period reached to 18.20% and 18.80% in the first and second seasons, respectively compared with the untreated ones. Therefore, this treatment could be recommended for raising garlic yield and improving bulb quality during the storage period under similar conditions to this work.

### **INTRODUCTION**

Garlic (*Allium sativum* L.) is one of the most important bulb vegetable crops. It plays dietary and medicinal roles in human being for countries. It has been cultivated since ancient times, used as a spice and flavoring. Because of its potential benefits in preventive and curative medicine, has been used in many cultures. (Rivlin, 2001). Even today, the medical use of garlic is widespread and growing (Amagase, 2006). In Egypt, it has been generally cultivated for both local consumption and export. Therefore, increasing garlic

yield and improving bulb quality are essential aims for both growers and consumers. Magnesium nutrition is one of major factors that affect growth, yield and quality of garlic. Its ions ( $Mg^{2+}$ ) have a specific role in the activation of enzymes involved in respiration, photosynthesis and the synthesis of DNA and RNA. Magnesium is also a part of the ring structure of the chlorophyll molecule. Studies indicate that 15 to 30% of the total magnesium in plants is associated with the chlorophyll molecule, the deficiency of magnesium will seriously affect of plant growth and development, being related directly to photosynthesis (Marschner, 1995).

The efficiency of fertilizers used in Egypt is low, either as a result of high pH of soil or high concentration of soil calcium carbonate. This problem could be solved by addition amounts of macro-elements fertilizers to the soil or through foliar application of them (Alexander, 1986). The positive effect of foliar application of macronutrients on growth, yield and chemical constituents of different plants may be attributed to the fact that these elements which can be readily absorbed by the leaves as a result of foliar spraying application and not lost through fixation, decomposition or leaching under unfavorable soils conditions (Doeing, 1986). Several attempts were done on the application of micronutrients spray to correct deficiency symptoms and enhance the vegetative growth of garlic which in turn reflects on increasing yield and its quality for facing local consumption and exportation (Eid *et al.*, 1991, Ibrahim *et al.*, 1991, Abdel-Fattah *et al.*, 2002, El-Morsy *et al.*, 2004 and El-Morsy, 2005). It can be, also used to satisfy acute needs of macro nutrients (Franke, 1986).

Several investigators indicated that spraying plants with magnesium enhanced plant growth, stimulated dry matter accumulation and increased yield and quality as well as chemical composition (Abd El-Rasoul and El-Azouni, 2002 on flax, Awad and El-Ghamry., 2007 on potato, Abo El-Hamd and Esmail, 2008 on sugar beet and Osman and El-Sawah, 2009 on tomato).

Yeast extract are the natural components (contains many compounds, i.e., cytokinins and proteins that enhance cell division and enlargement) which are safe and non-pollutant (Barnett *et al.*, 1990). Also, it contains the haloes-6-phosphate synthase (a key enzyme for trehalose biosynthesis) which not only affects plant development but also improves drought tolerance (Yeo *et al.*, 2000). Several investigators indicated that soaking cloves or spraying garlic plants enhanced plant growth, stimulated dry matter accumulation, increased bulb yield and quality and enhanced bulb storability (Tartoura and El-Saei., 2006, Abd El-Mageed *et al.*, 2009). Similar conclusions had been shown on other crops, i.e. Tomato growth and yield were increased by the foliar spraying with yeast extract (Fathy *et al.*, 2000 and Eata, 2001), also, foliar application of yeast extract significantly increased plant growth and yield of pea (Tartoura, 2001 and El-Desuki and El-Geready, 2006).

Thus, this study was planned to determine the effects of foliar application concentrations of magnesium as Mg-citrate and some yeast extract concentrations, in addition to their interactions on garlic productivity and storability under the conditions of Dakahlia Governorate.

## MATERIALS AND METHODS

Two field experiments were carried out in vegetable private Farm at Kafr Meet Faris, near El-Mansoura, Dakahlia Governorate, during two growing seasons of 2008/ 2009 and 2009/2010, to study the effect of some magnesium foliar application concentrations (0.0, 0.2%, 0.4% and 0.6% Mg as magnesium citrate 14.5% Mg) either single and/or in combination with some foliar application of yeast extract concentrations (0.0, 25 ml/L, 50 ml/L and 100 ml/L) on garlic (Sids-40) growth, yield and its components, as well as chemical constituents in cloves and bulb storability.

The experiment included 16 treatments which were 4 concentrations of magnesium and 4 concentrations of yeast extract as follows :

**a- Mg-concentrations:** Control treatment (spray only with water).

- 1- 0.2% magnesium.
- 2- 0.4% magnesium.
- 3- 0.6% magnesium.

**b- Yeast extract concentrations:**

- 1- Control treatment (spray only with water).
- 2- 25 ml/L.
- 3- 50 ml/L.
- 4- 100 ml/L.

Yeast extract was prepared according to procedure of Fathy *et al.* (2000) and Eata (2001), its chemical analysis according to methods of A. O. A. C. (1990).

Magnesium and yeast extract concentrations were supplied as a foliar application at 60, 75 and 90 days after planting. The control treatment was sprayed with tap water.

Garlic cloves were planted in the second week of October in both seasons. The experimental design was split plot with three replicates, the four foliar magnesium concentrations occupied the main plots which were subdivided to 4 sub plots each contained one of the yeast extract concentrations. Nearly uniform garlic cloves were soaked in running water for 12 h prior to planting and hand-planted at 10 cm apart on two sides of each row. All the plants were fertilized with the recommended doses of N, P and K. The other cultural practices for garlic commercial production were used according to the instructions laid down by the Ministry of Agriculture, Egypt. The harvesting time was in the second week of April, for both seasons.

**Data recorded:**

**Growth parameters:**

A random sample of five plants was taken from each plot after 120 days from planting to estimate plant height, number of leaves/plant, plant dry weight and bulbing ratio (neck diameter/bulb diameter).

**Yield and its components:**

At harvest time, marketable plants of each plot were cured, 15 days after harvest weighted in kg and converted to record as total yield (ton/fed). A random sample (10 bulbs) was taken from each treatment to determine bulb weight and diameter, as well as the number of cloves/bulb and clove weight .

**Chemical analysis:**

Samples of the dried cloves were ground, wet digested as described by Hesse (1971) and their nitrogen (N), phosphorus (P) and potassium (K) contents were determined according to the methods described by Chapman and Pratt (1961), John (1970) and Brown and Lilleland (1946), respectively. Total soluble solids (TSS) and volatile oils (mg/kg bulbs fresh weight) were determined according to A.O.A.C. (1990) and Gunther (1961).

**Storability:**

After curing, random samples (10 kg of marketable yield from every plot) were taken, stored at the normal room conditions (Table 1) and the percentage of weight loss was recorded monthly during the storage period (five months).

Data obtained during the two seasons of the study were statistically analyzed according to Gomez and Gomez (1984).

## RESULTS AND DISCUSSION

**Vegetative growth:**

Data presented in Table (1) show the effect of magnesium citrate concentrations, yeast extract concentrations and their interactions on growth aspects of garlic plants.

Concerning the effect of Mg-concentrations, it is clear from such data in Table (1) that plant height, number of leaves and plant dry weight in both seasons were significantly increased with increasing magnesium (Mg) concentration up to 0.6% Mg, also foliar application at this concentration was enhanced bulb:leaf ratio. These increases in growth parameters may be attributed to the effect of Mg on some physical functions such as carbohydrates synthesis and active many enzymes which in turn affect plant growth (Marschner, 1995). The obtained results concerted with those of (Abd El-Rasoul and El-Azouni, 2002 on flax, Awad and El-Ghamry., 2007 on potato, Abo El-Hamd and Esmail, 2008 on sugar beet and Osman and El-Sawah, 2009 on tomato).

Regarding, the effect of yeast extract concentrations, the same data in Table (1) reveal that foliar application of yeast extract concentrations had significant increases in all studied parameters of vegetative growth in both seasons. In this connection, plants sprayed with yeast extract at 50 ml/L were generally stocky and healthy in appearance than untreated plants. These results could be attributed to the great role of yeast in stimulate the cell division, elongation, enlargement, protein and nucleic acid synthesis and chlorophyll formation (Kraig and Haber., 1980, Spencer *et al.*, 1983). The obtained results are in harmony with those reported by Tartoura and El-Saei (2006) Abd El-Mageed *et al.* (2009). They mentioned that plant growth was enhanced with spraying yeast. Similarly, Abdel-Aziz (1997), mentioned that tomato growth parameters were responded to spray with yeast extract at 1 kg baker's yeast/200 L water.

**Table (1): Vegetative growth characters of garlic plants as affected by foliar Mg concentrations, Yeast and their interactions during 2008/2009 (S1) and 2009/2010 (S2) seasons.**

Characters Treatments	Plant height (cm)		Number of leaves/plant		Plant dry weight (gm)		Bulbing ratio		
	S1	S2	S1	S2	S1	S2	S1	S2	
<b>Mg-concentrations</b>									
Control	76.6	74.7	11.3	11.1	12.383	11.383	0.392	0.318	
0.2% Mg	78.1	76.8	11.6	11.6	12.033	12.100	0.392	0.312	
0.4% Mg	80.4	79.2	11.9	11.7	12.917	12.417	0.377	0.301	
0.6% Mg	82.2	81.8	12.3	11.8	13.700	13.317	0.368	0.348	
LSD at 5%	00.9	01.1	00.2	00.2	00.203	00.247	0.014	0.001	
<b>Yeast-concentrations</b>									
Control	77.0	75.8	11.4	11.0	12.000	11.700	0.397	0.371	
25 ml/L Y. extract	78.9	77.5	11.7	11.3	12.733	12.167	0.387	0.363	
50 ml/L Y. extract	81.8	80.0	12.3	12.1	13.700	13.900	0.367	0.342	
100 ml/L Y. extract	79.6	78.7	11.8	11.6	13.100	12.000	0.379	0.303	
LSD at 5%	00.7	00.6	00.2	00.3	00.265	00.261	0.001	0.001	
<b>Interactions:</b>									
<b>Mg-Concent.</b>		<b>Y.Concentrations</b>							
Control	Control	75.0	73.2	10.8	10.7	11.467	11.867	0.407	0.377
	25 ml/L	76.2	73.7	11.1	10.9	12.400	11.333	0.393	0.373
	50 ml/L	78.5	76.8	12.0	11.8	13.000	11.933	0.377	0.307
	100 ml/L	76.8	75.0	11.4	11.0	12.767	11.400	0.390	0.313
0.2% Mg	Control	75.8	73.8	11.2	11.0	11.767	11.733	0.403	0.377
	25 ml/L	77.2	76.3	11.4	11.4	12.267	12.000	0.397	0.367
	50 ml/L	80.7	79.2	12.1	12.1	13.333	12.700	0.380	0.347
	100 ml/L	78.8	77.8	11.6	11.7	12.867	12.267	0.390	0.307
0.4% Mg	Control	77.3	76.8	11.6	11.1	12.133	11.933	0.393	0.367
	25 ml/L	80.3	79.0	11.8	11.7	12.767	12.333	0.383	0.307
	50 ml/L	83.2	81.2	12.3	12.2	13.867	12.800	0.370	0.333
	100 ml/L	80.7	79.7	11.9	11.8	13.000	12.700	0.373	0.347
0.6% Mg	Control	79.8	79.3	11.9	11.3	12.933	12.467	0.383	0.363
	25 ml/L	81.8	80.8	12.2	11.4	13.200	13.000	0.373	0.307
	50 ml/L	85.0	83.0	12.8	12.4	14.700	14.267	0.303	0.330
	100 ml/L	82.0	82.3	12.3	11.9	13.867	13.733	0.363	0.343
L.S.D. at 5%		N.S	N.S	00.3	00.6	00.530	N.S	0.013	0.011

As for the interaction effects, it is obvious from the same data in Table (1) that all treatments of Mg-concentrations were generally more effective in the presence than in the absence of yeast extract. In this regard, plants sprayed with magnesium at 0.6% Mg and sprayed with yeast extract at 50 ml/L gave the highest values of plant growth in both seasons compared with the other treatments. Similar results were reported by Eata (2001) on tomato and Shokr and Fathy (2009) on snap bean.

**Yield and its components:**

Data illustrated in Table (2) show the effect of magnesium concentrations, yeast extract concentrations and their interactions on yield and its components of garlic. Such data indicate that foliar application of Mg at the high concentration (0.6% Mg) was generally beneficial than the other treatments. Moreover, this treatment significantly increased total yield, bulb weight and diameter as well as clove weight than the all studied Mg-concentrations in both seasons. However, number of cloves/bulb was

significantly affected by Mg-concentrations in the second season only. The positive effect of Mg-concentrations in improving total yield and its components may be attributed to the important role of Mg in increasing the activity of plant metabolism, which reflected on bulb yield and enhance bulb quality. In addition, the beneficial effect of Mg as a foliar fertilizer on the yield and its components may be due to the fact that Mg plays an important role in formation of the organic compound such as carbohydrates, lipids and etc...which translocate to the reproductive organs and consequently increasing the yield and its components (Marschner, 1995). These results are in agreement with those of (Abd El-Rasoul and El-Azouni, 2002 on flax, Hao-Xiuming and Papadopoulos, 2003 on tomato, Awad and El-Ghamry., 2007 on potato, Abo El-Hamd and Esmail, 2008 on sugar beet and Osman and El-Sawah, 2009 on tomato), they found that total yield, dray mattermarketable yield were increased with increasing Mg concentration.

**Table (2): Total yield and its components as affected by foliar Mg concentrations, Yeast and their interactions during 2008/2009 (S1) and 2009/2010 (S2) seasons.**

Characters	Total yield (ton/fed)		Bulb Weight (gm)		Bulb diameter (cm)		No. of cloves /bulb		Clove weight (gm)		
	S1	S2	S1	S2	S1	S2	S1	S2	S1	S2	
<b>Treatments</b>											
<b>Mg- concentrations</b>											
Control	0.930	0.778	02.070	09.783	0.9	0.7	10.0	10.7	3.182	2.962	
0.2% Mg	1.117	0.702	02.777	00.022	0.4	0.2	10.8	10.8	3.192	3.008	
0.4% Mg	1.797	0.980	08.722	02.070	0.5	0.2	10.1	10.9	3.787	3.097	
0.6% Mg	1.098	1.220	12.200	00.017	6.0	0.6	13.9	10.0	3.902	3.007	
LSD at 5%	0.252	0.135	02.192	01.193	0.2	0.2	N.S	00.6	0.304	0.160	
<b>Yeast-concentrations</b>											
Control	1.001	0.097	02.070	09.070	0.1	0.8	10.9	10.2	3.100	2.779	
25 ml/L Y. extract	1.201	0.780	00.822	00.700	0.4	0.1	10.0	10.1	3.202	2.901	
50 ml/L Y. extract	1.929	1.282	10.770	00.100	0.8	0.7	13.0	13.6	3.901	3.220	
100 ml/L Y. extract	1.718	1.028	08.002	02.882	0.0	0.3	10.3	10.0	3.709	3.207	
LSD at 5%	0.081	0.076	00.720	00.662	0.1	0.1	00.5	00.6	0.147	0.132	
<b>Interactions:</b>											
<b>Mg-conc. Y. Concentrations</b>											
Control	Control	0.070	0.208	08.867	07.967	0.4	0.2	10.2	10.7	2.722	2.767
	25 ml/L	0.098	0.022	09.122	07.900	0.0	0.8	10.7	10.7	2.900	2.800
	50 ml/L	1.397	1.102	07.100	02.022	0.3	0.0	13.2	13.7	3.782	3.000
	100 ml/L	1.177	0.787	00.200	00.722	0.0	0.9	10.2	10.7	3.270	2.980
0.2% Mg	Control	0.792	0.020	00.822	07.722	0.0	0.9	10.2	10.7	2.807	2.702
	25 ml/L	0.892	0.702	01.700	09.167	0.2	0.1	10.0	10.7	3.027	2.702
	50 ml/L	1.068	0.908	07.722	02.167	0.7	0.7	10.0	13.2	3.722	3.707
	100 ml/L	1.210	0.822	00.000	01.167	0.0	0.3	10.7	10.7	3.202	3.222
0.4% Mg	Control	1.200	0.700	00.722	09.022	0.3	0.0	10.7	10.2	3.207	2.770
	25 ml/L	1.000	0.928	07.067	02.067	0.0	0.2	10.2	10.7	3.027	3.087
	50 ml/L	1.102	1.282	12.767	00.100	0.8	0.7	13.2	10.2	0.177	3.002
	100 ml/L	1.802	1.062	10.067	02.200	0.0	0.3	10.0	10.2	3.787	3.082
0.6% Mg	Control	1.700	0.908	07.867	02.167	0.7	0.2	10.2	10.7	3.002	3.022
	25 ml/L	1.962	1.100	11.022	02.867	0.7	0.0	10.0	10.2	3.867	3.122
	50 ml/L	1.700	1.797	17.700	09.700	1.0	1.2	13.2	13.0	0.210	0.002
	100 ml/L	1.227	1.022	12.000	07.022	1.1	0.7	10.0	10.0	0.227	3.020
L.S.D. at 5%	N.S	0.152	01.441	01.352	0.2	N.S	01.0	01.2	0.294	N.S	

As for the effect of yeast extract concentrations, data in Table (2) indicate that total yield and its components were better with spraying the plants by yeast extract comparing with the untreated plants. Moreover, foliar application of yeast extract at 50 ml/L was more useful treatment to increasing total yield and improving its components than the other treatments. These increases might be ascribed to the fact that yeast contain sugar, proteins and amino acids, as well as several vitamins (Eata, 2001). The obtained results are in accordance with those of Tartoura and El-Saei (2006) Abd El-Mageed *et al.* (2009). Similar findings were established by El-Ghamriny *et al.* (1999) and Fathy *et al.* (2000) on tomato, Tartoura (2001) on Pea, El-Tohamy and El-Greadly (2007) on snap bean, Hanafy Ahmed, *et al.* (2007) on tomato.

Regarding the interaction effects, it is clear from data in Table (2) that the interactions between Mg-concentrations and yeast extract concentrations had a significant effect on total yield in the second season only while, bulb diameter and clove weight in the first season only, whereas, bulb weight and number of cloves/bulb were significantly affected in both seasons. In general, plants sprayed with 0.6% Mg concentration and 50 ml/L yeast extract produced the highest values. These results coincide with those of Shokr and Fathy (2009) on snap bean.

### **3- Chemical constituents:**

Data in Table (3) show the effect of magnesium concentrations, yeast extract concentrations and their interactions on element concentrations of N, P and K in cloves as well as percentage of total soluble solids and volatile oils in cloves of garlic.

From such data, it is evident that the Mg-concentrations had a significant effect on chemical constituents in cloves of garlic. All tested chemical constituents i.e., N%, P% and K% as well as TSS% and volatile oils were increased with increasing Mg-concentrations from 0 up to 0.6% Mg in both seasons. These increases in P and K percentages were significant in the first season only. These results are in agreement with those of Osman and El-Sawah (2009) on tomato.

Concerning the effect of yeast extract concentrations, data in Table (3) show that all concentrations of elements in cloves and percentage of total soluble solids and volatile oils in cloves were significantly increased due to plants sprayed with yeast extract compared with the untreated plants. The highest values of chemical concentration were produced by plants sprayed with 50 ml/L yeast extract in both seasons. These results agreed with those reported by Fathy *et al.* (2000) and Abou-Aly (2005) on tomato and El-Tohamy *et al.* (2007) on eggplant.

As for the interaction effects, it is evident from data in Table (3) that the interactions between Mg-concentrations and yeast extract concentrations had a significant effects on all concentrations of studied chemical constituents in cloves in both seasons. Plants sprayed with 0.6% Mg and 50 ml/L yeast extract achieved the highest concentrations of N, P and K, as well as TSS% and volatile oils.

**Table (3): Chemical constituents in garlic bulbs as affected by foliar Mg concentrations, Yeast and their interactions during 2008/2009 (S1) and 2009/2010 (S2) seasons.**

Characters	Macronutrients						TSS%		Volatile oils (Mg/100g f.w)		
	N %		P %		K %						
	S1	S2	S1	S2	S1	S2	S1	S2	S1	S2	
<b>Mg-concentrations</b>											
Control	1.33	1.30	0.37	0.40	1.31	1.33	2.21	2.23	0.378	0.393	
0.2% Mg	1.40	1.44	0.38	0.41	1.34	1.30	0.23	0.10	0.428	0.443	
0.4% Mg	1.46	1.48	0.44	0.46	1.36	1.38	0.40	0.70	0.448	0.467	
0.6% Mg	1.06	1.09	0.45	0.48	1.37	1.41	0.73	0.98	0.464	0.510	
LSD at 5%	0.01	0.01	0.01	N.S	0.01	N.S	0.06	0.10	0.004	0.008	
<b>Yeast-concentrations</b>											
Control	1.36	1.39	0.39	0.41	1.31	1.34	2.03	2.16	0.388	0.409	
25 ml/L Y. extract	1.41	1.44	0.40	0.43	1.33	1.30	2.18	0.1	0.418	0.444	
50 ml/L Y. extract	1.02	1.04	0.43	0.47	1.38	1.41	0.79	0.98	0.467	0.502	
100 ml/L Y. extract	1.46	1.49	0.42	0.44	1.30	1.38	0.27	0.02	0.446	0.463	
LSD at 5%	0.01	0.01	0.01	0.01	0.01	0.01	0.05	0.07	0.010	0.006	
<b>Interactions:</b>											
<b>Mg-Conc. Yeast-concentrations</b>											
Control	Control	1.20	1.27	0.30	0.37	1.28	1.31	3.73	3.87	0.307	0.373
	25 ml/L	1.30	1.33	0.34	0.38	1.29	1.32	3.90	4.10	0.373	0.390
	50 ml/L	1.38	1.41	0.41	0.43	1.34	1.36	4.90	0.20	0.407	0.420
	100 ml/L	1.37	1.39	0.38	0.40	1.33	1.34	4.30	4.07	0.377	0.390
0.2% Mg	Control	1.33	1.36	0.30	0.37	1.29	1.32	4.30	4.03	0.380	0.400
	25 ml/L	1.36	1.38	0.37	0.39	1.34	1.34	4.73	4.80	0.420	0.437
	50 ml/L	1.01	1.03	0.41	0.44	1.37	1.39	0.87	0.83	0.460	0.480
0.4% Mg	Control	1.38	1.41	0.42	0.44	1.34	1.30	4.80	0.13	0.393	0.417
	25 ml/L	1.42	1.40	0.43	0.40	1.30	1.37	0.10	0.43	0.430	0.407
	50 ml/L	1.07	1.08	0.40	0.49	1.39	1.41	7.10	7.40	0.490	0.523
	100 ml/L	1.47	1.46	0.44	0.47	1.36	1.40	0.70	0.83	0.477	0.473
0.6% Mg	Control	1.49	1.02	0.44	0.46	1.34	1.37	0.30	0.00	0.420	0.447
	25 ml/L	1.00	1.08	0.44	0.47	1.30	1.38	0.47	0.70	0.400	0.493
	50 ml/L	1.72	1.70	0.46	0.50	1.43	1.47	7.30	7.00	0.510	0.587
L.S.D. at 5%	0.01	0.01	0.01	0.01	0.01	0.01	0.10	0.15	0.013	0.011	

**4- Storability:**

Data in Table (4) show the effect of magnesium concentrations, yeast extract concentrations and their interactions on monthly weight loss percentage of garlic bulbs during the storage period. Such data indicate that foliar application of Mg at the high concentration (0.6% Mg) was generally beneficial than the other treatments. The weight loss percentages during and at the end of storage period were decreased with increasing the foliar application concentration up to 6% Mg-citrate in both seasons. These results may be due to increase dry matter in plants (Table 1), TSS % and chemical constituents in cloves (Table 3).

**Table (4): Weight loss percentage of garlic as affected by foliar Mg concentrations, yeast and their interactions during 2008/2009 (S1) and 2009/2010 (S2) seasons.**

Characters	Weight loss (%) during the storage period										
	30 days		60 days		90 days		120 days		150 days		
	S1	S2	S1	S2	S1	S2	S1	S2	S1	S2	
<b>Mg-concentrations</b>											
Control	22.7	22.5	27.0	27.8	21.4	23.0	27.7	27.9	29.7	20.5	
0.2% Mg	22.1	22.7	26.7	27.1	20.9	23.0	27.0	27.0	28.7	29.3	
0.4% Mg	22.8	28.7	21.0	22.3	22.5	28.2	28.0	20.7	21.0	23.0	
0.6% Mg	21.3	28.0	29.5	21.0	21.7	20.3	22.6	27.9	27.3	29.0	
LSD at 5%	01.1	01.1	01.4	0.6	01.2	01.0	01.5	01.5	00.5	00.2	
<b>Yeast- concentrations</b>											
Control	21.8	22.0	20.4	26.7	29.7	22.3	22.5	27.6	27.2	29.3	
25 ml/L Y. extract	20.1	21.1	22.2	20.5	27.7	20.7	22.7	23.7	20.1	27.0	
50 ml/L Y. extract	22.9	29.1	22.0	22.8	22.8	27.3	29.2	29.6	21.7	27.6	
100 ml/L Y. extract	29.1	20.1	22.2	22.4	26.2	29.2	21.0	21.7	22.3	22.5	
LSD at 5%	00.4	00.6	00.4	00.6	00.4	00.6	00.3	00.6	00.5	00.2	
<b>Interactions:</b>											
<b>Mg-concent.</b>	<b>Yeast-Concent.</b>										
Control	Control	22.8	22.0	27.7	28.2	22.3	22.2	29.0	21.2	22.2	22.0
	25 ml/L	22.7	22.7	27.1	28.0	21.9	22.9	28.7	28.1	20.3	21.0
	50 ml/L	22.0	22.9	27.2	27.2	20.2	21.6	20.8	20.8	27.2	28.1
	100 ml/L	22.0	22.2	27.9	27.7	20.9	22.2	27.2	27.0	29.1	28.9
0.2% Mg	Control	22.8	22.5	27.1	27.8	22.2	22.7	28.2	28.1	20.3	21.1
	25 ml/L	22.2	22.9	27.0	27.5	21.2	22.9	27.1	27.6	29.2	20.1
	50 ml/L	21.1	21.9	26.0	26.2	29.1	22.1	22.7	22.8	27.0	27.0
	100 ml/L	22.2	22.2	27.8	27.9	20.8	22.1	27.0	20.9	28.0	29.0
0.4% Mg	Control	21.9	22.2	22.2	26.2	28.2	22.1	22.2	20.2	20.2	28.0
	25 ml/L	28.1	28.8	22.2	22.2	20.2	29.6	29.1	21.2	22.2	22.0
	50 ml/L	20.0	26.1	29.0	20.0	21.0	22.0	20.2	27.0	28.2	20.0
	100 ml/L	26.2	27.2	20.6	22.8	22.0	27.1	27.2	29.0	20.0	22.0
0.6% Mg	Control	28.7	20.2	22.2	22.2	20.2	29.1	28.2	21.8	21.0	22.9
	25 ml/L	22.2	29.2	20.5	22.1	22.1	27.6	20.6	28.7	28.1	21.0
	50 ml/L	22.7	20.2	26.6	27.7	28.8	21.2	21.2	22.0	22.0	20.2
	100 ml/L	20.2	27.1	28.6	20.1	20.6	22.0	22.2	20.0	27.1	28.0
L.S.D. at 5%	00.8	01.2	00.7	01.3	00.8	01.2	00.6	01.2	01.1	00.4	

Regarding the effect of yeast extract concentrations, the same data in Table (4) indicate that bulb storability of plants sprayed with yeast extract was better than that of the untreated plants. Moreover, foliar application of yeast extract at 50 ml/L was more beneficial than the application once. These results are in harmony with those of Tartoura and El-Saei (2006) and Abd El-Mageed *et al.* (2009) they found that weight loss percent of bulbs was significantly reduced during the storage period with plants sprayed by yeast extract.

Concerning the interaction between foliar spray of magnesium and yeast extract concentrations, data in Table (4) show that the positive interactions often observed on storability of bulbs. The lowest total weight loss percentages during and at the end of the storage period were obtained from foliar spray with 0.6% Mg and 50 ml/L yeast extract in both seasons. From the obtained results of this study, it could be concluded that, spraying garlic plants with combination between 0.6% Mg and 50 ml/L yeast extract

was the superior treatment to enhancing the garlic plant growth, yield and its components, as well as bulb quality and storability. This treatment achieved increases in yield at the end of the storage period reached to 18.20% and 18.80% in the first and second seasons, respectively compared with the untreated ones. Therefore, this treatment could be recommended under similar conditions to this work.

## REFERENCES

- A. O. A. C. (1990). Official methods of analysis. 15<sup>th</sup> ed. A.O.A.C., Washington, Dc, USA.
- Abd El-Mageed, I.Y. T.; Youssef, N. S. and Shereen M. Youssef (2009). Effect of some anti oxidant sources on growth, yield and storability of garlic. *Egypt J. Agric. Res.*, 87 (1): 245-264.
- Abd El-Rasoul, SH. M. and A.M.A. El-Azouni (2002). Effect of magnesium, farmyard manure and mineral fertilizers on yield, yield components and nutrient contents of some flax genotypes on sandy soils. *J. Agric. Sci. Mansoura Univ.*, 27 (11): 7951-7962.
- Abdel-Aziz, M.A. (1997). Response of tomato plants to nitrogen fertilizer concentrations and growth regulators. M.Sc.Thesis, Dept.of Vegetable Crops, Faculty of Agric.Cairo Univ.Egypt.
- Abdel-Fattah, A.E.; Z.S. El-Shal and E.M. Awad (2002). Response of garlic productivity and storability to application times of phosphorus and certain micro-nutrients.*J.Agric. Sci. Mansoura Univ.*, 27 (9): 6231-6241.
- Abo El-Hamd, A.S. and A.A. Esmail (2008). Effect of foliar potassium, magnesium and boron on yield and quality of sugar beet (*beta vulgaris*, L.)growing in new reclaimed soils. *J. Agric. Sci. Mansoura Univ.*, 33 (3): 1737-1746.
- Abou-Aly, H.E. (2005). Stimulatory effect of some yeast applications on response of tomato plants to inoculation with biofertilizers. *Annals of Agricultural Science, Moshtohor*, 43(2):595-609.
- Alexander, A. (1986). Foliar fertilization. ed. By Alexander. Schering Agrochemical Division, Special fertilizer group, Berlin (FRG), P. 17-60.
- Amagase, H. (2006). Clarifying the real bioactive constituents of garlic, *Journal of Nutrition*, 131, 7165-7255.
- Awad, El. M. M. and A. M. El-Ghamry (2007). Effect of humic acid, effective microorganisms (em) and magnesium on potatoes in clayey soil. *J. Agric. Sci. Mansoura Univ.*, 32 (9): 7629-7639.
- Barnett, J.A., Payne, R.W., and Yarrow, D. (1990). Yeasts, characteristics and identification. Cambridge Univ. Press Publ. By the Press Syndicate of the Univ. of Cambridge Camb. PP. 999.
- Brown, T. D. and O. Lilleland (1946). Rapid determination of potassium and sodium in plant materials and soil extracts by flame photometry. *Proc. Amer. Soc. Hort. Sci.*, 48: 301-304.
- Chapman, H. D. and P. F. Pratt (1961). Methods of analysis for soil, plant and water. Soil dept. and plant nutrition, California Univ., Citrus Exp. Sta. Riverside, California.

- Doeing, H.W. (1986). Foliar fertilization. An effect method for controlling micronutrients deficiencies on calcareous and saline soil in arid lands. In: Proc. Symp. Application of Special Fertilizers. Alex. 21-23 Feb., Egypt (Ed.).
- Eata, A.M. (2001). Response of some tomato cultivars to natural soil salinity and use of some treatments to reduce salt injury. Ph. D Thesis, Fac. Agric. Mansoura Univ., Mansoura, Egypt.
- Eid, S.M.; N.S. Shafshak and F.A. Abo-Sedera (1991). Effect of potassium fertilization and foliar application of certain micro-nutrients combinations on growth, yield and chemical composition of garlic plants. *Annals Agric. Sci. Moshtohor*, 29 (2): 981-993.
- El-desuki, M., and Nadia H.M. El-Geready (2006). Response of pea plants to foliar application of yeast extract. *J. Agric. Sci. Mansoura Univ.*, 31 (10): 6667-6674.
- El-Ghamriny, E.A., Arisha, H.M.E. and K.A. Nour (1999). Studies on tomato flowering, fruit set, yield and quality in summer season. 1- Spraying with thiamine, ascorbic acid and yeast. *Zagazig J.Agric.Res.*, 26 (5):1345-1364.
- El-Morsy, A.H.A. (2005). Effect of sulphur concentrations and foliar application of certain micronutrients on garlic (*Allium sativum* L.). The Sixth Arabian Horticulture Conference, Ismailia, Egypt, March 20-22, 457-467.
- El-Morsy, A.H.A.; Z.S. El-Shal and Sawsan M. H. Sarg (2004). Effect of potassium application methods and some micronutrients on growth, yield and storability of garlic. *J. Agric. Sci. Mansoura Univ.*, 29 (4): 2013-2023.
- El-Tohamy, W.A. and N.H.M. El-Greadly (2007). Physiological responses, growth, yield and quality of snap beans in response to foliar application of yeast, vitamin E and zinc under sandy soil conditions. *Australian Journal of Basic and Applied Sciences*, 1(3): 294-299.
- Fathy, El-S.L.El-S, Farid, S. and S.A. El-Desouky (2000). Induce cold tolerance of outdoor tomatoes during early summer season by using triphosphate (ATP), yeast, other natural and chemical treatments to improve their fruiting and yield. *J. Agric. Sci. Mansoura Univ.*, 25 (1):377-401.
- Franke, W. (1986). The basis of foliar absorption of fertilizers with special regard to the mechanisms. "Foliar fertilization" ed. By Alexander. Schering Agrochemical Division, Special fertilizer group, Berlin (FRG), P. 17-25.
- Gomez, K.A. and A.A. Gomez (1984). "Statistical Procedure for agric. Res." 2<sup>nd</sup>. Ed. John-Wiley and Sons. Inc. New Yourk, 680 pages.
- Gunther, E. (1961). "The Essential Oils" 4<sup>th</sup>. Ed. Vol. 1 D., van Nostrand Co. Inc., New York.
- Hanafy Ahmed, A.H., Khalil, M.K., Metwally, A.M. and M.A. Abdel-Aziz (2007). Effect of biomagical, yeast, acadian sea plants, foliafeed d, trial compound fertilizer, putrescine and silicon treatments on growth, yield and chemical composition of tomato plants grown under early and late summer seasons. *J.Agric. Sci.Mansoura Univ.*, 32 (2):913-949 .

- Hao-XiuMing and A.P. Papadopoulos (2003). Effects of calcium and magnesium on growth, fruit yield and quality in a fall greenhouse tomato crop grown on rockwool. *Canadian Journal of Plant Science*, 83(4): 903-912.
- Hesse, P.R. (1971). A text book of soil chemical analysis. John-Murray (Puplish), London, Great Britain.
- Ibrahim, D.M.; A.R. Ahmed and A.M. Abd El-Hameid (1991). Influence of zinc and manganese fertilization treatments on bulb components and chemical composition of Balady and Chinese garlic cultivars. *Zagazig J. Agric. Res.*, 18 (2): 511-519.
- Jackson, M.L. (1967). Soil chemical analysis. Prentice-Hall of India, Newdelhi, India. P. 115.
- John, M.K. (1970). Colorimetric determination of phosphorus in soil and plant material with ascorbic acid. *Soil Sci.*, 109 : 214-220.
- Kraig, E. and J.E. Haber (1980). Messenger ribonucleic acid and protein metabolism during sporulation of *Saccharomyces cervisiae*. *J. Bacterial*, 144: 1098-1112.
- Marschner, H. (1995). Mineral Nutrition of Higher Plants, 2<sup>nd</sup> Ed. Academic Press, Inc. San Diego, CA, USA, 889 pp.
- Osman, A. Sh. And Nevein A. El-Sawah (2009). Response of tomato plants grown under reclaimed soil conditions to foliar application of magnesium. *J. Agric. Sci. Mansoura Univ.*, 34 (5): 4929-4942.
- Rivlin, R. (2001). Historical perspective on the use of garlic. *Journal of Nutrition*, 136, 9515-9545.
- Shokr, M. M. B. and El-S. L. El-S. Fathy (2009). Some foliar applications for improving snap bean (*phaseolus vulgaris*, l.) Quality and yield at fall season. *J. Agric. Sci. Mansoura Univ.*, 34(5): 5089-5106.
- Spencer, T.F.T.; S.M. Dorothy and A.R.W. Smith (1983). Yeast genetics fundamental and applied aspects. Pp. 16-18. ISBN0-387-90973-9. Springer-Verlag New York, USA.
- Tartoura, E.A.A. and M. A. El-Saei (2006). Effect of compost tea, yeast extract and their mixture on yield and storability period in garlic. *J. Agric. Sci. Mansoura Univ.*, 31 (7): 4801-4811.
- Tartoura, E.A.A (2001). Response of pea plants to yeast extract and two sources of n-fertilizers. *J. Agric. Sci. Mansoura Univ.*, 26 (12): 7887-7901.
- Yeo, E., HawkBin, K., SangEun, H., JoonTak, S. E., JinChang, R., MyungOk, B., Yeo, E. T., Kwon, H. B., Han, S. E., Lee, J. T., Ryu, J. C. and M. O. Buyn (2000). Genetic engineering of drought resistant potato plants by introduction of the trehalose-6-phosphate synthase (TPSI) gene from *Saccharomyces cerevisiae*. *Molecules and Cells*. 10(3) 263-268.

## استجابة نمو وإنتاجية وقابلية تخزين الثوم للرش بالماغنسيوم ومستخلص الخميرة.

عبد الله حلمي على المرسي ، أسامة محمد سيف الدين و عبد البديع صالح عزت  
قسم بحوث الخضر - معهد بحوث البساتين - مركز البحوث الزراعية - الجيزة - مصر.

تعتبر إضافة العناصر المغذية للنباتات بالرش الورقي أكثر سرعة وكفاءة لسد احتياج النبات من العناصر الصغرى عنها بالإضافة الأرضية، ويمكن أيضاً الاستفادة من هذه الطريقة لسد احتياجات النبات من بعض العناصر الكبرى.

ولهذا الغرض، نُفذت تجربتان حقليتان على محصول الثوم (صنف سدس-٤٠) في مزرعة خضر خاصة بكفر ميت فارس بالقرب من المنصورة بمحافظة الدقهلية خلال موسم الزراعة ٢٠٠٨/٢٠٠٩ و ٢٠٠٩ / ٢٠١٠ م لدراسة تأثير بعض تركيبات الرش الورقي بعنصر الماغنسيوم (٠.٢%، ٠.٤%، ٠.٦% ماغنسيوم في صورة سترات الماغنسيوم) كلٌّ منها منفرداً أو مع الرش بمستويات مختلفة من مستخلص الخميرة (٢٥ مل/لتر، ٥٠ مل/لتر و ١٠٠ مل/لتر بالإضافة لمعاملة الكنترول) على نمو النباتات ومحصول الأصيل ومكوناته بالإضافة إلى المحتويات الكيماوية في الفصوص والقدرة التخزينية للأبصال خلال فترة التخزين (٥ شهور). وقد وزعت المعاملات في قطع منشقة مرة واحدة في ثلاثة مكررات، ويمكن تلخيص النتائج المتحصل عليها فيما يلي :-

بصف عامة أوضحت النتائج أن الرش بالماغنسيوم عند تركيز ٠.٦% إلى حدوث زيادات ملموسة في ارتفاع النبات، عدد الأوراق و الوزن الجاف لعرش النبات كما تحسنت نسبة التبصيل وكذلك ازداد المحصول الكلي ومتوسط وزن وقطر البصلة وبجانب ذلك زادت معنوياً تركيبات المواد الصلبة الكلية والزيوت الطيارة وكذلك زادت نسبة عناصر النيتروجين والفوسفور والبوتاسيوم في فصوص الثوم، وقد أدت أيضاً إلى حدوث انخفاض معنوي في نسبة نقص وزن الأصيل خلال فترة التخزين في كلا موسمي الدراسة.

ومن ناحية أخرى أدى رش النباتات بمستخلص الخميرة بتركيز ٥٠ مل/لتر إلى حدوث زيادات معنوية في معظم صفات النمو الخضري للنباتات وكذلك المحصول الكلي ومكوناته، كما أدى إلى زيادة تركيبات المواد الصلبة الكلية والزيوت الطيارة والنيتروجين والفوسفور في الفصوص مقارنة مع معاملة الكنترول، وبجانب ذلك أدى إلى انخفاض نسبة نقص وزن الأصيل عند نهاية فترة التخزين معنوياً.

التفاعلات بين تركيبات الرش بالماغنسيوم و تركيبات الرش بمستخلص الخميرة لوحظت في حالات كثيرة ، ولقد كانت أفضل النتائج باستخدام الرش الورقي بالماغنسيوم عن تركيز ٠.٦% مع الرش بمستخلص الخميرة عند تركيز ٥٠ مل/لتر، كما أدت هذه المعاملة إلى زيادة في المحصول في نهاية فترة التخزين تُقدر بـ ١٨.٢٠% و ١٨.٨٠% في الموسم الأول والموسم الثاني على التوالي مقارنة بمعاملة الكنترول.

وبناءً على ماتقدم، يمكن التوصية باستخدام هذه المعاملة لرفع إنتاجية الثوم وتحسين جودة الأصيل وقابليتها للتخزين تحت الظروف المشابهة لظروف هذا البحث.

قام بتحكيم البحث

كلية الزراعة - جامعة المنصورة  
مركز البحوث الزراعية

أ.د / سمير طه محمود العفيفي  
أ.د / عبد المنعم محمد عبد الحميد