

Using of Tamarind and Vanilla Essential Oils for the Management of Brown Rot Disease of Potato

Moussa, Z.¹; Eman Z. Goma²; Ehsan M. M. Rashad³ and E. A. Salem⁴

¹Bacterial Diseases Research Dept., Plant Pathology Research Institute, Agricultural Research Center, Giza, Egypt.

²Biological and Geological Science Dept., Faculty of Education, Ain Shams University, Cairo, Egypt.

³Microbiology Activity Unit, Microbiology Dept., Soils, Water and Environment Research Institute, Agricultural Research Center, Giza, Egypt,

⁴Food Irradiation Dept., National Center for Radiation Research and Technology, Atomic Energy Authority, Cairo, Egypt

Corresponding author: Moussa, Z.: Bacterial Diseases Research Dept., Plant Pathology Research Institute, Agricultural Research Center, Giza, Egypt – e.mail: zeadmoussa@gmail.com



ABSTRACT

The hazard of chemical pesticides on the environment is a worldwide problem. Finding ecofriendly alternatives to chemical pesticides is a recent trend in agricultural research, one of the best choices is essential oils (EOs). This study aims to manage brown rot disease of potato using EOs. *In vitro*, Seven of 16 EOs inhibited *Ralstonia solanacearum*. The minimal inhibitory concentration (MIC) value was determined on *R. solanacearum* growth, tamarind and vanilla EOs had the lowest MIC (1%). Gas Chromatography-Mass Spectrometry analysis indicated that the main components of tamarind EO (TEO) and vanilla EO (VEO) are propylene glycol (50.97 %) and vanillin (40.78 %), respectively. TEO had a stronger bactericidal effect than VEO when *R. solanacearum* incubated in a solution of MIC of both EOs. In the pots experiment, soaking potato slices (spunta cultivar) in 1% of each of TEO and VEO for 30 minutes significantly improved peroxidase, polyphenol oxidase, chlorophylls and carotenoids, as well as, increased the plant height after 70 days of planting in comparison with the control treatment. Both EOs significantly decreased the disease rating from 4.2 in control treatment to 1.6 (TEO) and 2.2 (VEO). In addition to marked increments up to 36.11% (TEO) and 32.52% (VEO) in fresh tubers weight. After 60 days of tubers storage at room temperature, the two EOs decreased the infected tubers by 69.05% (TEO) and 47.17% (VEO). Generally, TEO showed better results than VEO. The growth inhibition of *R. solanacearum* and enhancement of physiological characters and yield of potato, encouraging the use of these two uncommon EOs as effective tools to manage brown rot disease of potato.

Keywords: *Ralstonia solanacearum*, Inhibition, Bactericidal effect, Pot Experiment, Yield, Physiological characters.

INTRODUCTION

Brown rot (or bacterial wilt) disease of potato is one of the world's most devastating disease observed in most potato-producing areas of the world. The causal bacterial pathogen is called *Ralstonia solanacearum* that has a wide host range, it can infect plants belonging to 54 families; more than 450 plant species. It has high persistence and unavailable resistant crop varieties (Patil *et al.*, 2017).

The chemical pesticides cause pollution that leads to hazards for the environment and human beings. So, researchers try to find safe ecofriendly tools to fight different plant pest and diseases. Essential oils are tools from these tools (Pavela and Benelli, 2016). Different methods of control of brown rot disease of potato, such as physical, cultural, biological, chemical, and integrated management were investigated (Yuliar *et al.*, 2015). Essential oils (EOs) are promising environmentally friendly tools in the management of *R. solanacearum* (Vu *et al.*, 2017).

The EOs are complex, volatile, secondary metabolites. These natural compounds are synthesized by different parts of aromatic plants. They have inhibitory effect against bacteria, fungi, (Miller *et al.* 2015) and viruses (Sánchez, and Aznar 2015) Furthermore, they have insecticidal (Pavela and Benelli, 2016) and nematocidal effect (Avato *et al.*, 2017), in addition to the pharmaceutical, medicinal and cosmetic applications (Bakkali *et al.*, 2008).

The EOs were used as effective tools in the management of soil born plant pathogen (Mihajlović *et al.*, 2017). Moreover, the EOs are considered a new horizon to fight the bacterial antibiotic resistance (Yap *et al.*, 2014).

Oboo *et al.* (2014) revealed that the EOs of *Tarchonanthus camphorates* (Camphor brush), *Ocimum suave* (Cambodia) and *Lippie javanica* (Sage brush) significantly reduced the bacterial wilt disease of potato.

Li and Yu (2015) reported that the EO obtained from the leaves of *Macleaya cordata* R. Br. inhibited the growth of *R. solanacearum*. They showed that their EO increased the permeability of *R. solanacearum* cell membrane causing leakage of electrolytes, the losses of reducing sugar and proteins and change in the cell morphology leading to the death of bacteria.

The EOs enhance the physiological characters and increase the systemic induced resistance of plants and therefore increase the plant defense against pathogens. The EOs increase induced resistance of plants against bacterial pathogens (Lucas *et al.*, 2012) and fungal pathogens (Vergnes *et al.*, 2014).

The chemical components of the EOs vary from plant to another. Moreover, they vary in the same plant according to the effect of different factors that include the soil conditions, seasonal variation (humidity, temperature, rain, etc.), the age of the plant, the part of the plant and the method of extraction (Evergetis *et al.*, 2016). The EOs are very complex natural compounds that contain different volatile components at different concentrations. They have two or three main components at concentrations (20 to 70%) in comparing with others existing in trace concentrations. The EOs have different biological effects. The different components of the EOs have a synergistic effect. However, it is probable that the activity of the main components is modified by other minor molecules (Bakkali *et al.*, 2008).

Pino *et al.* (2004) found that the major volatile constituents of tamarind (*Tamarindus indica* L.) fruits were 2-phenylacetaldehyde, 2-furfural and hexadecenoic acid. While Escalona-Arranz *et al.* (2010) reported that the leaves extract of tamarind had antimicrobial activity and the main components of this extract were benzyl benzoate, limonene and hexadecanole.

The main volatile components of vanilla (*Vanilla planifolia*) were vanillic acid (4%–5%), 4-hydroxy - benzaldehyde (6% – 9%) and vanillin (85%–87%) (Takahashi *et al.*, 2013). Vanilla EO (VEO) inhibited a soil-borne Gram-negative bacterium that is called *Chromobacterium violaceum* CV026 where it inhibited quorum-sensing genes expression of the Tn-5 mutant (Choo *et al.*, 2006). Also, *in vitro*, VEO had inhibitory effects against *Escherichia coli*, *Streptococcus faecalis*, *Enterobacter aerogenes*, *Proteus aeruginosa* and *P. vulgaris* (Subramanian *et al.*, 2009).

This investigation aims to use some uncommon EOs in control of brown rot disease of potato. The Effect of the EOs on inhibition of *R. solanacearum*, as well as, minimal inhibitory concentration (MIC) values of the effective ones were investigated. The bactericidal effect of the most effective EOs on *R. solanacearum* was studied. Pots experiment was carried out to indicate the *in vivo* effectivity of the most effective EOs on the management of the studied disease and on potato plants.

MATERIALS AND METHODS

Essential oils:

The used 16 commercial EOs were obtained from Kheder El-Attar Academy, Cairo, Egypt. The tested EOs were: Tamarind (*Tamarindus indica* L.), Vanilla (*Vanilla planifolia*), Hyacinth (*Hyacinthus orientalis*), Prunus (*Prunus domestica*), Apple (*Malus domestica*), Jasmine (*Jasminum sambac*), Pineapple (*Ananas comosus*), Tea (*Camellia sinensis*), Thermos (*Lupinus albus* L.), Olive (*Olea europaea* L.), Visnaga (*Ammi visnaga* (L.) Lam.), Limon (*Citrus × limon* (L.) Burm.f.), Banana (*Musa* sp.), Juniper (*Juniperus* L.), Flax (*Linum usitatissimum* L.) and Castor (*Ricinus communis*).

Bacterial pathogen:

Virulent isolate of *Ralstonia solanacearum* (race 3 biovar 2) was obtained from the Bacterial Collection of Bacterial Diseases Research Department, Plant Pathology Research Institute, Agricultural Research Center, Giza, Egypt

Potato Tubers: Potato tubers (Spunta cultivar) was obtained from Nubaria district, Bahaira Government, Egypt.

Antibacterial activity:

The agar disc diffusion method was carried out according to Teixeira *et al.* (2013) to study the antibacterial effect of the EOs against *R. solanacearum*. A bacterial suspension of *R. solanacearum* was adjusted to 10^8 CFU/mL and spread onto nutrient glucose agar (NGA) plates. Plates were kept for drying. On the center of plate, sterilized filter paper discs (5 mm; Whatman No. 1), that drenched with the individual EO, were placed. Negative controls were prepared with sterilized distilled water. Plates were staying at 4 °C for 2h for permission the diffusion of EOs from the discs through the media. The incubation of these plates was carried out at 28 °C for 48 h. The inhibitory effect of tested EOs was assessed by measuring the radius of inhibition zone (if present). Three replicates were prepared for each tested EO.

Minimal inhibitory concentration (MIC) determination:

EOs that gave the largest inhibition zones were selected to investigate their MICs. The MIC is the

lowest concentration which inhibits microorganism's growth. The MICs for the selected EOs were determined by using agar dilution method (Balouiri *et al.*, 2016). A serial dilution of each EO (2, 1, 0.5 and 0.25 % (v/v), was prepared in molten NGA medium with 0.5% (v/v), Tween-80 was incorporated into the agar after autoclaving. Inoculation was carried out with streaking *R. solanacearum* by a sterilized loop from liquid culture (10^8 CFU/ml 48 h. old) onto the prepared plates. Plates of NGA medium, with 0.5% (v/v) Tween-80 and without any EO, was used as a control treatment. The incubation of these plates was carried out for 48 h. at 28 °C. The EOs having the lowest MICs were selected for further studies.

Analysis of the selected essential oils:

The selected EO(s) were subjected to analysis by Gas Chromatography/ Mass Spectrometry (GC/MS) which was performed in Central Agricultural Pesticide Laboratory (CAPL), Dokki, Giza, Egypt. GC/MS was performed on an Agilent 6890 gas chromatograph equipped with an Agilent mass spectrometric detector, with a direct capillary interface and fused silica capillary column PAS-5 ms (30 mm X 0.25 um film thickness). The injection of samples was performed under the following conditions:

The carrier gas washes helium (approximately 1 ml/min., pulsed splitless mode). The injection size was 1.0 ul. The solvent delay was 3 min. The mass spectrophotometric detector was operated in electron impact ionization mode an ion energy of 70 e.v. scanning from m/z 50 to 500. The ion source temperature was 230 °C and the quadrupole1 temperature was 150 °C. The electron multiplier voltage (EM voltage) was maintained 1250v above autotuned. The instrument was manually tuned using perfluorotributyl amine (PFTBA). The GC temperature program was started at 50 °C for 3 min. then elevated to 280 °C at a rate of 8 °C/min. and 15 min. hold at 280 °C the injector temperature was set at 280 °C. Willy and Nist 05 mass spectral data base was used in the identification of the separated peaks.

Solubility test:

The solubility of the MIC of the selected EOs in water (without tween 80) was tested.

Bactericidal effect of the selected EOs:

The bactericidal effect of the selected EOs was investigated according to the method of Joray *et al.* (2011). The suspension of *R. solanacearum* was prepared with concentration about 10^{10} CFU/ ml. The control treatment was prepared by adding 10 ml of the prepared bacterial suspension to 90 ml of sterilized tap water (or with Tween 80). The treatments of selected EOs were prepared as aforementioned with control and adding each individual selected EO with concentration of its MIC. In an incubator shaker (28 °C and 120 rpm/min), the flasks were incubated. At selected time intervals, the serial dilution was applied for samples of these treatments and plated onto NGA plates. The incubation of plates was performed at 28 °C for 48 h. After incubation, colony forming units (CFU) were determined. The values of log CFU/ml were calculated. Five replicates were applied for each dilution.

Pots experiment:

The preparation of inoculum of bacterial pathogen (*R. solanacearum*) was done by inoculating

this pathogen in Nutrient Glucose Broth (NGB) and incubating in an incubator shaker for 48 h. (28 °C and 120 rpm). By centrifugation (10000 rpm/ min for 20 min.), bacteria were precipitated. In sterilized tap water, the precipitate was re-suspended. The bacterial suspension was adjusted to 10⁸ CFU/ ml concentration.

This experiment was carried out during the period of 21 February 2016 to 31 May 2016 at an open-air at Tag El-Ezz Agricultural Research Station, Dakahlia, Egypt. The plastic sacks (pots) [35 cm diameter] were filled with 20 kg non-sterilized soil. The used soil was a mixture of clay and sand 1:1 (w: w). Each pot was irrigated by water. Pots were left to dry for 72 h., then infestation was done by 100 ml of *R. solanacearum* (10⁸ CFU/ ml) for each pot. For about 48 h. the pots were left to dry. The healthy potato tuber slices of similar size were soaked (for 30 min.) in the selected EOs with concentration equal MIC of each selected EO. In each pot, two potato tuber slices were planted. After 40 days of planting, one plant from each pot was harvested for assessment of polyphenol oxidase and peroxidase enzymes in addition to total phenols. The other plant of each pot was harvested at the end of the experiment. The plant height was measured after 70 days of planting. In addition, for estimation of chlorophylls and carotenoids, leaves from the same locations in the stem of potato plants were taken. The disease rating was estimated. For each treatment, five replicates were prepared. The experiment was designed in a completely randomized block design.

Biochemical assessments:

The enzymatic activity of peroxidase and polyphenol oxidase were assessed by using a spectrophotometric method according to Seleim *et al.* (2104). Folin-Ciocalteu reagent method [as described by Blainski *et al.* (2013)] was used to estimate total phenolic contents of fresh leaves. Chlorophyll a, chlorophyll b, total chlorophyll, and carotenoids in potato leaves were estimated as described by Wellburn (1994).

Disease rating:

As the scale described by Kempe and Sequeira (1983), wilt symptoms were assessed, where, (0 = no symptoms, 1= up to 25 % wilt, 2 = 26-50 % wilt, 3 = 51-75 % wilt, 4 = 76-99% of the foliage wilted and 5 = dead plants. Each plant was examined, the disease rating was assessed, and the mean value for each treatment was calculated.

Tubers weight:

At the harvesting time, the tubers of each pot were weighted. The mean weight and the percentage in the increase of tubers weight were calculated.

The percentage of the increase of potato tubers weight = [(Tubers weight of treatment – tubers weight of control) / tubers weight of control] X100

Infected tubers percentage:

Potato tubers were kept at room temperature for 6 weeks to help the appearance of latent infection, then, symptoms were clearly detected (Graham *et al.*, 1979). Potato tubers were cut – after the storage period - to examine the existence of the disease symptoms. The infected tubers number was recorded. The percentage of the infected tubers was determined. The decrease of the percentage of infected tubers was calculated as the following:

The decrease in infected tubers % = [(Infected tubers % of control - Infected tubers % of treatment)/ infected tubers % of control] X100

Statistical analysis:

statistical analysis by software packages CoStat (version 6.4, CoHort Software, U.S.A) was used. The one-way randomized blocks design was applied in pots experiment. Duncan’s multiple range test was used to compare the means at probability (P) value of ≤ 0.05.

RESULTS AND DISCUSSION

Results in (Table 1) showed that from 16 EOs, seven EOs were effective in inhibition of the growth of *R. solanacearum* *in vitro* experiment. The other tested EOs did not effective against the growth of *R. solanacearum*. Tamarind EO (TEO) was the most effective one causing inhibition zone of 40 mm diameter. Vanilla EO (VEO) caused inhibition zone of 36 mm diameter. On the other hand, jasmine EO had the smallest inhibition zone (11 mm diameter). This finding had the same harmony of the results of Li and Yu (2015) who found - *in vitro* - that the EO obtained from the leaves of *Macleaya Cordata* (Willd.) R. Br. inhibited *R. solanacearum*.

Also, results in Table (1) revealed the MIC values of TEO and VEO were 1%. Hyacinth EO had MIC value 2 %. Prunus and apple EOs had MIC values more than 2 %. Therefore, TEO and VEO were selected for further investigations in this study. These results had the confirmation of those of Escalona-Arranz *et al.* (2010) who found that tamarind extract from leaves had antimicrobial activity. Choo *et al.* (2006) indicated that VEO inhibited quorum-sensing genes expression of the Tn-5 mutant of *Chromobacterium violaceum* CV026.

Table 1. The inhibitory effect of the tested EOs against *R. Solanacearum* and minimal inhibitory concentration of most effective oils.

No.	Essential Oil	Inhibition zone diameter (mm) *	Minimal inhibitory concentration (MIC)
1	Tamarind	40	1.0 %
2	Vanilla	36	1.0 %
3	Hyacinth	30	2%
4	Prunus	30	> 2%
5	Apple	25	> 2%
6	Pineapple	21	N.D.
7	Jasmine	11	N.D.
8	Tea	0	N.D.
9	Thermos	0	N.D.
10	Olive	0	N.D.
11	Visnaga	0	N.D.
12	Limon	0	N.D.
13	Banana	0	N.D.
14	Juniper	0	N.D.
15	Flax	0	N.D.
16	Castor	0	N.D.

* The inhibition zone diameters were measured directly from the plate, the recorded numbers were the means of three replicates, 0: No inhibition zone and N.D.: Not detected.

Data in Table (2) indicated the GC/MS of TEO and VEO. Data revealed that propylene glycol is the main component (50.97 %) of TEO and other components are Benzaldehyde (39.70%), 4-[(dimethylamino) methyl]-2-methoxyphenol (6.92%) and 2 (3H)-Furanone (1.77 %). The

main component of VEO is vanillin (40.78 %) and the other components are 2(3H)-furanone (21.75 %), 3-ethoxy-4-hydroxy-Benzaldehyde (19.24 %), acetic acid (10.43 %), benzyl alcohol, (5.26 %) and Vanillin propylene glycol acetal (2.24 %). GC/MS analysis of TEO disagrees with the results of volatile components of TEO obtained by Pino *et al.* (2004) and Escalona-Arranz *et al.* (2010). Li and Yu (2015) found that the chemical composition of EO of their study is different from the chemical composition of the same EO of the same plant. The differences of volatile components may be due to several reasons, such as, the part of the plant used for extraction, physiological age of the plant, method of extraction, environmental condition and the growing season (Li and Yu, 2015).

Propylene glycol has bactericidal activity on the growth of *S. mutans*, *E. faecalis* and *E. coli* (Nalawade *et al.*, 2015). Moreover, “2 (3H)-Furanone” have antimicrobial activity against six types of bacteria and two types of fungi *in vitro* (Abou-Elmagd *et al.*, 2015). Also, benzaldehyde has insecticidal, antimicrobial, and antioxidant activity (Ullah *et al.*, 2015).

Vanillin has antibacterial activity against *Listeria innocua*, *Lactobacillus plantarum* and *Escherichia coli* acting as a membrane-active compound, resulting in the dissipation of ion gradients and the inhibition of respiration, the extent to which is species-specific. These effects initially do not stop the production of ATP Fitzgerald *et al.* (2004). Acetic acid has antimicrobial activity against different isolates of *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Streptococcus pyogenes* and *Proteus mirabilis* (Thorp *et al.*, 1998). Moreover, it inhibited the growth of *Shigella* species: *S. dysenteriae*, *S. sonnei*, *S. boydii* and *S. flexneri*, (In *et al.*, 2013). Bjarnsholt *et al.* (2015) showed that acetic acid has antibiofilm properties.

Table 2. Gas Chromatography/ Mass Spectrometry (GC/MS) of TEO and VEO.

Essential oil	Component	Area %
TEO	Propylene glycol	50.97
	2 (3H)-Furanone	1.77
	Benzaldehyde	39.70
	4-[(dimethylamino)methyl]-2-methoxyphenol	6.92
VEO	Acetic acid	10.43
	Benzyl alcohol	5.26
	2(3H)-Furanone	21.75
	Vanillin	40.78
	3-ethoxy-4-hydroxy-Benzaldehyde	19.24
	Vanillin propylene glycol acetal	2.24

The test of the solubility of the two selected EOs in water without tween 80 indicated that these two EOs are soluble in water without tween 80. Therefore, the following tests were carried out by dissolving these two EOs directly in water without tween 80. Therefore, this solubility in water enhanced the antibacterial activity of these two EOs (Calo *et al.*, 2015).

As shown in Fig. (1) the incubation of *R. solanacearum* with TEO and VEO at concentration of 1 % (MIC) led to significant decrements of bacterial count of *R. solanacearum* more than the control (incubation with sterilized tap water). TEO was more effective in

decreasing the viable bacterial count of *R. solanacearum* than VEO. The decrease in the viable bacterial count of *R. solanacearum* that preserved in sterilized tap water is very low; Log CFU/ml decreased from 9.876 to 9.140 after 24 h. The reduction in the count of *R. solanacearum* that incubated with TEO was high till 3 h. where the Log CFU/ml was decreased from 9.162 to 3.788. While, after 3 h., the decrease of the bacterial count is very low, where Log CFU/ml was decreased from 3.788 after 3 h. of incubation to 3.352 after 24 h. of incubation. A similar trend was observed with VEO where the rate of decreasing of the bacterial count was high till 3 h. and after 3 h. the rate of decreasing of bacterial count till 24 h. is very low. These results are in the same trend of Li and Yu (2015) who found that the count of *R. solanacearum* count was decreased when it incubated with the MIC of the EO from the leaves of *Macleaya cordata* R. Br. The mode of action of the two selected EOs against *R. solanacearum* may be due increasing the permeability of bacterial cell membrane leading to leakage of electrolytes, the losses of reducing sugar and proteins and damage in the bacterial cells leading to death of bacteria and decreasing of the viable bacterial count according to the findings of Li and Yu (2015).




Fig. 1. Bactericidal effect of the two selected EOs on *R. solanacearum*

All recorded values are means of five replicates.

Data in Table (3) declared that the two selected EOs improved the productivity of enzymes that play an important role in inducing resistance and photo-synthetic pigments. The two selected EOs led to significant increase in peroxidase (POD) activity, which increased from 12.00 Unit.min⁻¹.g⁻¹ FW in control to 22.67 and 16 Unit.min⁻¹.g⁻¹ FW in TEO and VEO treatments, respectively. The polyphenol oxidase (PPO) increased from 5.33 Unit.min⁻¹.g⁻¹ in control treatment to 13.33 and 6.67 Unit.min⁻¹.g⁻¹ in the plants treated with TEO and VEO respectively where the last increase was non-significant. Furthermore, total phenols increased in plants treated with the two EOs from 27.84 mg GAE/g FW to 53.89 and 38.53 mg GAE/g FW in TEO and VEO treatments, respectively. These physiological parameters reflecting the health condition of the plant. These results are confirmed with Vergnes *et al.* (2014) who found that the EOs increase induced resistance against the bacterial pathogen or the fungal pathogen. In this respect, Lamba *et al.* (2008) reported phenols accumulated rapidly of at the infection site, which limited or slowed the growth of the pathogen due to their antimicrobial and antioxidant action. The proposed mode of action for the pathogen defense role of PPO, include; (1) General toxicity to pathogens. (2) Alkylation causing reduction of bioavailability of cellular proteins to the pathogen. (3) A physical barrier to pathogens

in the cell wall that is forming due to crosslinking of quinones with protein or other phenolics. (4) Quinone redox cycling, leading to H₂O₂ and other reactive oxygen species that are important factors in plant pathogen interactions and defense signaling (Raj *et al.*, 2006). POD is an important factor in the integrated resistance response of plants to different of stresses due to the its important roles in the production of toxic secondary metabolites, in cell wall toughening and its simultaneous oxidant and antioxidant capabilities (Thakker *et al.*, 2013).

The two EOs treatments led to significant increases in the photosynthetic pigments of potato leaves compared with the control (Table 3). Potato plants treated with TEO

recorded the highest values in the content of Chl a, b and total Chl (1.225, 0.8923 and 2.0908 mg/g FW, respectively), followed by VEO (1.0720, 0.8658 and 1.9643 mg/g FW, respectively). The same trend was observed in carotenoids. Enhancement of the chlorophyll content is a parameter indicating improving the health condition of the plant, by enhancing the effectiveness of photosynthetic apparatus leading to enhancement in the disease resistance (Amaresh and Bhatt, 1998). Carotenoids are essential for plant life, during photosynthesis. Carotenoids are parts of a framework of protection of the photosynthetic apparatus against high light disorder (Lachman *et al.*, 2016).

Table 3. Physiological characteristics of potato as affected by the two EOs treatments in pots experiment

Treatment	Peroxidase (Unit.min ⁻¹ . g ⁻¹ FW)	Polyphenol oxidase (Unit.min ⁻¹ . g ⁻¹ FW)	Total phenols (mgGAE/g FW)*	Photo-synthetic pigments (mg/g FW)			
				Chlorophyll a	Chlorophyll b	Total Chlorophyll	Carotenoids
Control	12.00 c	5.33 b	27.84 c	0.9792 c	0.8100 c	1.7892 b	0.5269 b
Tamarind oil	22.67 a	13.33 a	53.89 a	1.2250 a	0.8923 a	2.0908 a	0.6317 a
Vanilla oil	16.00 b	6.67 b	38.53 b	1.0720 b	0.8658 b	1.9643 a	0.6609 a

*: milligram Gallic Acid equivalent (GAE)/ gram Fresh weight.

All recorded values are means of three replicates; means followed by the different letter within each column is significantly different using Duncan's Multiple Range Test at P value of ≤0.05

As revealed in Fig. (2), the two selected EOs enhanced the growth of potato plants. The plant height after 70 days of planting was significantly increased in plants treated with EOs comparing with control. The plant height increased from 31.2 cm in the control to 43.2 and 41 cm in the plants treated with TEO and VEO, respectively. This improvement of vegetable growth may be due to the enhancement of physiological characters and the inhibitory effect of two EOs against the tested bacterial pathogen.




Fig. 2. Effect of the two selected EOs on plant height of potato pants after 70 days of planting.

Column scripted with different letter is significantly different using Duncan's Multiple Range Test at P value of ≤0.05.

Data illustrated in Fig. (3) indicated that disease rating of treated plants with the two selected EOs was significantly decreased than the corresponding value of the untreated plants. The disease rating decreased from 4.2 in the control to 1.6 and 2.2 in plants treated with TEO and VEO, respectively. These results had the confirmation of Oboo *et al.*, (2014) who indicated that EOs decreased the disease incidence of bacterial wilt of potato. The decrease of disease rating may due to the inhibitory effect of the two EOs against the test bacterial pathogen and the increase of systemic induced resistance of potato plant.

Data illustrated in Fig. (4) showed that the tubers weight of the treated pots significantly increased compared with the control. The weight of tubers for each pot increased from 122.4 g in the control to 166.6

and 162.2 g in the plants treated with TEO and VEO respectively. The percentages of yield increase were 36.11 % and 32.52 % with TEO and VEO, respectively.




Fig. 3. Disease rating of potato plants in pots experiment

Column scripted with different letter is significantly different using Duncan's Multiple Range Test at P value of ≤0.05. Disease rating scale: 0 = no symptoms, 1= up to 25 % wilt, 2 = 26-50 % wilt, 3 = 51-75 % wilt, 4 = 76-99% of the foliage wilted and 5 = dead plants




Fig. 4. Effect of the two selected EOs on tubers weights of potato tubers.

Column scripted with different letter is significantly different using Duncan's Multiple Range Test at P value of ≤0.05.

Data illustrated in Fig. (5) revealed that the infected tubers percentage of the two treatments of the used EOs after 60 days of storage at room temperature were significantly decreased. These percentage decreased from 88.33 % in the control treatment to 27.33 and 46.33 % with TEO and VEO treatment, respectively. While there was no significant difference between the value of this parameter with the treatments of the two EOs. So, the treatment of potato seeds

with these two EOs decreased the infected tubers by percentage 69.05 % for TEO and 47.17 for VEO. The increase in yield and decrease in potato infected tubers percentage may be because of these two EOs against the studied bacterial pathogen and their effect on other bacterial and fungal pathogens (Miller *et al.*,2016) and their insecticidal effect (Pavela and Benelli, 2016). EOs have nematocidal activity (Avato *et al.* 2017). In addition, this effect is a result of their enhancement of physiological characters of the plant and increase in the induced resistance.




Fig. 5. Effect of the two selected EOs on the infected tubers percentage after storage for 60 days at room temperature.

Column scripted with different letter is significantly different using Duncan's Multiple Range Test at P value of ≤ 0.05

CONCLUSION

Therefore, it could be concluded that TEO and VEO can be used to manage brown rot disease of potato. *In vitro*, they inhibited the growth of *R. solanacearum*. In pots experiment, they significantly decreased the disease rating and increased plant height during the vegetative growth. They enhanced the potato plants systemic induced resistance and improve their physiological characters. They significantly increased the fresh weight of potato tubers and significantly decreased the infected potato tubers.

REFERENCES

Abou-Elmagd, W. S., El-Ziaty, A. K. and A. A. Abdalha 2015. Ring transformation and antimicrobial activity of indolyl-substituted 2 (3H)-furanones. *Heterocyclic Communications*, 21(3), 179-185.

Amaresh, C. and R.K. Bhatt 1998. Biochemical and physiological response to salicylic acid in reaction to systemic acquired resistance. *Photosynthetica*, 35: 255-258.

Avato, P., Laquale, S., Argentieri, M. P., Lamiri, A., Radicci, V. and T. D'Addabbo, 2017. Nematicidal activity of essential oils from aromatic plants of Morocco. *J. Pest Sci.*, 90 (2), 711-722.

Bakkali, F.; Averbeck, S.; Averbeck, D., and M. Idaomar 2008. Biological effects of essential oils—a review. *Food and Chem. Toxicol.* 46 (2), 446-475.

Balouiri, M., Sadiki, M., and S. K. Ibensouda 2016. Methods for *in vitro* evaluating antimicrobial activity: A review. *J. Pharm. Analysis*, 6 (2), 71-79.

Bjamsholt, T., Alhede, M., Jensen, P. Ø., Nielsen, A. K., Johansen, H. K., Homøe, P., Højby N., Givskov, M. and K. Kirketerp-Møller, 2015. Antibiofilm properties of acetic acid. *Advances in Wound Care*, 4 (7), 363-372.

Blainski, A., Lopes, G. C., and J. C. P. De Mello 2013. Application and analysis of the Folin Ciocalteu method for the determination of the total phenolic content from *Limonium brasiliense* L. *Molecules*, 18: 6852-6865.

Calo, J. R., Crandall, P. G., O'Bryan, C. A., and S. C. Ricke 2015. Essential oils as antimicrobials in food systems – A review. *Food Control*, 54, 111-119.

Choo, J.H.; Rukayadi, Y. and J.K. Hwang 2006. Inhibition of bacterial quorum sensing by vanilla extract. *Lett. Appl. Microbiol.*, 42, 637-641.

Escalona-Arranz, J. C.; Péres-Roses, R.; Urdaneta-Laffita, I.; Camacho-Pozo, M. I.; Rodríguez-Amado, J.; and I. Licea-Jiménez 2010. Antimicrobial activity of extracts from *Tamarindus indica* L. leaves. *Pharmacognosy Magazine*, 6 (23), 242 - 247.

Evergetis, E., Michaelakis, A., Papachristos, D. P., Badieritakis, E., Kapsaski-Kanelli, V. N., and S. A. Haroutounian 2016. Seasonal variation and bioactivity of the essential oils of two *Juniperus* species against *Aedes* (*Stegomyia*) *albopictus* (Skuse, 1894). *Parasitol. Res.*, 115 (6), 2175-2183.

Fitzgerald, D.J.; Stratford, M.; Gasson, M.J.; Ueckert, J.; Bos, A. and A. Narbad 2004. Mode of antimicrobial action of vanillin against *Escherichia coli*, *Lactobacillus plantarum* and *Listeria innocua*. *J. A. Microbiol.*, 97, 104-113.

Graham, J., Jones, D.A. and A.B. Loyd, 1979. Survival of *Pseudomonas solanacearum* race 3 in the Plant debris and in latent infected potato tubers. *Phytopathol.*, 69:1100 – 1103.

In, Y. W., Kim, J. J., Kim, H. J. and S. W. Oh 2013. Antimicrobial activities of acetic acid, citric acid and lactic acid against *Shigella* species. *J. Food Safety*, 33(1), 79-85.

Joray, M.B.; Del Rollan, M.R.; Ruiz, G.M.; Palcios, S.M. and M.C. Carpinella 2011. Antibacterial activity of extracts from plants of central Argentina – isolation of an active principle from *Achrocline satuireioides*. *Planta Med.* 77, 95 – 100.

Kempe, J. and L. Sequeira 1983. Biological control of bacterial wilt of potatoes: Attempts to induce resistance by treating tubers with bacteria. *Plant dis.*, 67(5): 499-503.

Lachman, J., Hamouz, K., Orsák, M., and Z. Kotíková, 2016. Carotenoids in potatoes – a short overview. *Plant Soil Environ.*, 62:10, 474-481.

Lamba, P., Sharma, S., Munshi, G.D. and S.K. Munshi 2008. Biochemical changes in sunflower plants due to seed treatment/spray application with biocontrol agents. *Phytoparasitica*, 36: 388-399.

Li, C. M. and J. P. Yu 2015. Chemical composition, antimicrobial activity and mechanism of action of essential oil from the leaves of *Macleaya Cordata* (Willd.) R. Br. *J. Food Safety*, 35(2), 227-236.

Lucas, G. C., Alves, E., Pereira, R. B., Perina, F. J., and R. M. D. Souza 2012. Antibacterial activity of essential oils on *Xanthomonas vesicatoria* and control of bacterial spot in tomato. *Pesquisa Agropecuária Brasileira*, 47(3), 351-359.

Mihajlović, M.; Rekanović, E.; Hrustić, J., Grahovac, M. and B. Tanović 2017. Methods for management of soilborne plant pathogens. *Pesticidi I Fitomedicina*, 32(1), 9-24.

- Miller, A. B., Cates, R. G., Lawrence, M., Soria, J. A. F., Espinoza, L. V., Martinez, J. V. and D. A. Arbizú (2015). The antibacterial and antifungal activity of essential oils extracted from Guatemalan medicinal plants. *Pharm. Biol.*, 53(4), 548-554.
- Nalawade, T. M., Bhat, K. and S.H. Sogi 2015. Bactericidal activity of propylene glycol, glycerine, polyethylene glycol 400, and polyethylene glycol 1000 against selected microorganisms. *J. Int. Soc. Prev. Community Dent.* 5 (2):114-9.
- Oboon, H., Muia, A. W., and Z. M. Kinyua 2014. Effect of selected essential oil plants on bacterial wilt disease development in potatoes. *J. App. Biosci.*, 78 (1), 6666-6674.
- Patil, V. U., V. Girimalla, V. Sagar, R. S. Chauhan and S. K. Chakrabarti. 2017. Genome sequencing of four strains of phytotypes I, II and IV of *Ralstonia solanacearum* that cause potato bacterial wilt in India. *Barazil. J. Microbiol.*, 48, 193-195.
- Pavela, R., and G. Benelli 2016. Essential oils as ecofriendly biopesticides? Challenges and constraints. *Trends Plant Sci.*, 21(12), 1000-1007
- Pino, J. A.; Marbot, R. and C. Vazquez 2004. Volatile components of tamarind (*Tamarindus indica* L.) grown in Cuba. *J. Essential Oil Res.*, 16(4), 318-320.
- Raj, S. N., Sarosh, B. R., and H. S. Shetty 2006. Induction and accumulation of polyphenol oxidase activities as implicated in development of resistance against pearl millet downy mildew disease. *Functional Plant Biol.*, 33(6), 563-571.
- Sánchez, G., and R. Aznar 2015. Evaluation of natural compounds of plant origin for inactivation of enteric viruses. *Food Environ. Virol.*, 7(2), 183-187.
- Seleim, M. A., Abo-Elyousr, K. A., Mohamed, A. A. A. and H. A. Al-Marzoky, 2014. Peroxidase and polyphenoloxidase activities as biochemical markers for biocontrol efficacy in the control of tomato bacterial wilt. *J. Plant Physiol. Pathol.* 2: 1-4.
- Subramanian, S.; Banu, H.H.; Bai, R.M.R. and R. Shanmugavalli 2009. Biochemical evaluation of antihyperglycemic and antioxidant nature of *Psidium guajava* leaves extract in streptozotocin-induced experimental diabetes in rats. *Pharm. Biol.*, 47, 298-303.
- Takahashi, M.; Inai, Y.; Miyazawa, N.; Kurobayashi, Y. and A. Fujita 2013. Identification of the Key Odorants in Tahitian Cured Vanilla Beans (*Vanilla tahitensis*) by GC-MS and an Aroma Extract Dilution Analysis. *Biosci. Biotechnol. Biochem.*, 77, 601-605.
- Teixeira, B., Marques, A., Ramos, C., Neng, N. R., Nogueira, J. M., Saraiva, J. A., and M. L. Nunes 2013. Chemical composition and antibacterial and antioxidant properties of commercial essential oils. *Industrial Crops and Products*, 43, 587-595.
- Thakker, J. N., Patel, S., and P. C. Dhandhukia 2013. Induction of defense-related enzymes in banana plants: Effect of live and dead pathogenic strain of *Fusarium oxysporum* f. sp. cubense. *ISRN Biotechnology*, 1-6.
- Thorpe, M., Kruger, J., Oliver, S., Nilssen, E., and C. Prescott 1998. The antibacterial activity of acetic acid and Burrow's solution as topical otological preparations. *J. Laryngol. Otol.*, 112(10), 925-928.
- Ullah, I., Khan, A. L., Ali, L., Khan, A. R., Waqas, M., Hussain, J., Lee, J.H. and J. H. Shin 2015. Benzaldehyde as an insecticidal, antimicrobial, and antioxidant compound produced by *Photobacterium temperata* M1021. *J. Microbiol.*, 53(2), 127-133.
- Vergnes, S., Ladouce, N., Fournier, S., Ferhout, H., Attia, F., and B. Dumas 2014. Foliar treatments with *Gaultheria procumbens* essential oil induce defense responses and resistance against a fungal pathogen in Arabidopsis. *Frontiers Plant Sci.*, 5: 1-8.
- Vu, T. T., Choi, G. J., and J. C. Kim 2017. Plant-derived Antibacterial Metabolites Suppressing Tomato Bacterial Wilt Caused by *Ralstonia solanacearum*. *Res. Plant Dis.*, 23(2), 89-98.
- Wellburn, A. R. (1994). The spectral determination of chlorophylls a and b, as well as total carotenoids, using various solvents with spectrophotometers of different resolution. *J. Plant Physiol.*, 144(3), 307-313.
- Yap, P. S. X., Yiap, B. C., Ping, H. C., and S. H. E. Lim 2014. Essential oils, a new horizon in combating bacterial antibiotic resistance. *Open Microbiol. J.*, 8, 6-14.
- Yuliar, Nion, Y. A., and K. Toyota 2015. Recent trends in control methods for bacterial wilt diseases caused by *Ralstonia solanacearum*. *Microbes and Environments*, 30(1), 1-11.

استخدام الزيوت العطرية للتمر الهندي والفانيليا لمكافحة مرض العفن البني للبطاطس

زيداد موسى^١، إيمان زكريا جمعة^٢، إحسان محمد محمد رشاد^٣ و إيهاب أحمد سالم^٤
^١قسم بحوث الأمراض البكتيرية - معهد بحوث أمراض النباتات - مركز البحوث الزراعية - الجيزة - مصر
^٢قسم العلوم البيولوجية والجولوجية - كلية التربية - جامعة عين شمس - القاهرة - مصر.
^٣وحدة النشاط الميكروبي - قسم الميكروبيولوجي - معهد بحوث الأراضي والمياه والبيئة - مركز البحوث الزراعية - الجيزة - مصر.
^٤قسم تشعب الأغذية - مركز بحوث وتكنولوجيا الإشعاع - هيئة الطاقة الذرية - القاهرة - مصر.

تمثل أضرار المبيدات الكيميائية على البيئة مشكلة عالمية، والعثور على بدائل صديقة للبيئة لمبيدات الآفات الكيميائية هو اتجاه حديث في البحوث الزراعية، وتعد الزيوت العطرية واحد من أفضل هذه الخيارات. تهدف هذه الدراسة إلى مكافحة مرض العفن البني في البطاطس باستخدام الزيوت العطرية. ومن بين ١٦ من الزيوت العطرية، سبعة منها ثبتت نمو بكتيريا رالستونيا سولاناسيرم في المعمل. ولقد تم تحديد أقل تركيز مثبط على نمو البكتيريا، وسجل زيتي التمر الهندي والفانيليا أقل تركيز مثبط (١٪). ولقد بين تحليل الغاز الكروماتوجرافي والطيف الكتلي أن المكونات الرئيسية لزيتي التمر الهندي والفانيليا هي البروبيلين جليكول (٩٧.٥٠٪) والفانيلين (٧٨.٤٠٪) على التوالي. وكان لزيت التمر الهندي تأثير مبيد على البكتيريا أقوى من زيت الفانيليا وذلك عندما تم اختبارهما على البكتيريا في محلول يحتوي أقل تركيز مثبط من كلا الزيتين. وفي تجربة الأوص أدى غمر شرائح البطاطس (صنف سبونت) في ١٪ من كل من الزيتين لمدة ٣٠ دقيقة إلى تحسن معنوي في البيروكسيداز، البوليفينول أو أكسيديز والكلوروفيل والكاروتينات، وكذلك زيادة ارتفاع النبات بعد ٧٠ يوما من الزراعة بالمقارنة مع معاملة المقارنة. وقد أدت المعاملة بكل الزيتين العطريين إلى انخفاض معنوي في مرض العفن البني من ٤.٢ في معاملة الكنترول إلى ١.٦ (معاملة التمر الهندي) و ٢.٢ (معاملة الفانيليا) بالإضافة إلى زيادات ملحوظة تصل إلى ٣٦.١١٪ (التمر الهندي) و ٣٢.٥٢٪ (الفانيليا) في وزن الدرناط الطازجة. وعند تخزين الدرناط لمدة ٦٠ يوما عند درجة حرارة الغرفة، خفضت المعاملة بكل الزيتين العطريين من الدرناط المصابة بنسبة ٦٠.٦٪ (التمر الهندي) و ٤٧.١٧٪ (الفانيليا). عموما، أظهر زيت التمر الهندي نتائج أفضل من زيت الفانيليا في تثبيط نمو بكتيريا رالستونيا سولاناسيرم وتحسن في الخواص الفسيولوجية وإنتاجية البطاطس، مما يشجع على استخدام هذين الزيتين العطريين الغير شائعين كأدوات فعالة لمكافحة مرض العفن البني في البطاطس.