

**CHEMICAL COMPOSITION, ADULTICIDAL AND REPELLENT ACTIVITY OF ESSENTIAL OILS FROM *Mentha longifolia* L. AND *Lavandula dentata* L. AGAINST *Culex pipiens* L.**

**Ali, S. Al-Sarar**

Plant Protection Department, College of Food and Agriculture Sciences, King Saud University, P.O. Box 2460, Riyadh 11451, Saudi Arabia.

**ABSTRACT**

The essential oils from *Mentha longifolia* L., and *Lavandula dentata* L. were evaluated for their insecticidal and repellent activity against adult females of *Culex pipiens* L. The two oils showed LC<sub>50</sub> values of 0.215 and 0.217 µl / l, respectively. *Lavandula dentata* oil showed higher repellent activity than *Mentha longifolia* oil against adults of *Culex pipiens*; paraffin oil significantly prolonged the time of protection for the two oils. The longest time of protection was 165 min for *L. dentata* oil, at 1 µl/cm<sup>2</sup>, when the tested oils were applied in paraffin oil. The principle compounds in the two oils were pulegone, l-menthone, 1-8-cineole, eucarvone and borneol for (*M. longifolia*); and camphor, fenchone, linalool and fenchol for (*L. dentata*). The two oils were effective as repellent substances when compared with commercial materials. Further studies are needed to isolate the most effective mosquito control agents from these oils.

**Keywords:** essential oils, *Culex pipiens*, adulticidal activity, *Mentha longifolia*, repellents, *Lavandula dentata*

**INTRODUCTION**

Mosquitoes are serious pests; they annoy man and animals and transmit many fatal diseases (Service, 2004; Smith *et al.*, 1993; Sardelis, 2003; Peng *et al.*, 2004; Morales *et al.*, 2006). Conventional pesticides are heavily used to combat mosquitoes, but these pesticides have many disadvantages as: development of resistance, residue problems and bad effects on environment and human beings (WHO, 1992; Briassoulis, 2001). Therefore, a need is found to replace these hazardous pesticides with safe ones. Plant-derived pesticides are being tried as optional alternatives. While a number of botanicals, especially essential oils, were reported to be effective as mosquito repellents (Sritabutra and Soonwera, 2013; Phasomkusolsil and Soonwera, 2011; Gillij *et al.*, 2008; Prajapati *et al.*, 2005), many exhibited larvicidal activity (Govindarajan *et al.*, 2013; Manimaran *et al.*, 2012; Ansari *et al.*, 2000; Cetin *et al.*, 2004; Cetin and yanikoglu, 2006; Michaelakis *et al.*, 2007). Few studies were carried out on adulticidal properties of essential oils against mosquitoes (Pal *et al.*, 2011; Kang *et al.*, 2009; Dolan, *et al.*, 2007; Yang *et al.*, 2005). At least, five-seven years are required to introduce a new commercial insecticide into the market; therefore, it is important to discover the pesticidal properties of local plant species to be used as soon as the onset of insect resistance to conventional pesticides. The two medicinal

plants, *Mentha longifolia* L., and *Lavandula dentata* L. were reported to have insecticidal and/or antimicrobial properties in many places (Odeyemi *et al.*, 2008; Imelouane *et al.*, 2009; Koliopoulos *et al.*, 2010; Hafedh *et al.*, 2010; Hussain *et al.*, 2010; Baka, 2010). Therefore, this work was undertaken to investigate the constituents of *Mentha longifolia*, and *Lavandula dentata* oils and evaluate their insecticidal and repellent activity against *Culex pipiens* adults.

## **MATERIALS AND METHODS**

### **The tested Plants:**

The samples of *L. dentata* were collected from Asir region, Saudi Arabia, while samples of *M. longifolia* were purchased from a local supermarket. The tested Plants were identified by the Botany Department, King Saud University.

### **Extraction of essential oils**

Semi-dried leaves and flowers of *L. dentata* and fresh leaves of *M. longifolia* were steam distilled. Essential oils were extracted from steam distillates with diethyl ether; extracts were dried over anhydrous sodium sulfate and solvent was evaporated under vacuum, using Rotavapor-R-215 rotary evaporator. Oils were kept in deep freezer, in glass vials with Teflon screw caps.

### **GC-MS analysis**

The components of the tested oils were identified by Agilent Technologies 6890 N Network system, equipped with Agilent 5973 mass selective detector. Agilent DB-35 MS capillary column was used with the following oven temperature program: 40°C initial temperature, 2 °C /min ramp to 200 °C, held for 5 min. Carrier gas (helium) flow rate was 1 ml/min, injection of oil sample (1µl dissolved in methanol) was carried out at 200 °C. Electron ionization mode with a transfer line temperature at 280 °C was used. A Wiley 7 data base was employed to identify the compounds.

### **Insecticidal activity**

#### **Insects tested**

Adult females of *Culex pipiens* were obtained from a susceptible strain reared in the insectary of Plant Protection Department, King Saud University. The colony was kept at 24 ± 2 °C, with a photoperiod of 12: 12 h (light:dark). Larvae were fed on animal feed and adults were fed on 10% sucrose.

#### **Adulticidal trials**

The fumigant toxicity of the tested oils against female *C. pipiens* was performed in airtight conical flasks according to our method (Al-Sarrar *et al.*, 2014), with minor modifications. The tested oil was pipetted into a 3 ml glass vial. The opening of the treated vial was covered with a cloth netting to prevent contact of mosquitoes with the oil inside the vial. Twenty adult females (5-10-days-old) were introduced into the conical flask with the aid of an aspirator. The treated vial was hanged inside the conical flask with a string. The conical flask was sealed with its quick fit lid. To prevent any leakage of oil vapor outside the flask, the lid was wrapped with parafilm.

Dead insects were counted 24 h after treatment. Overturned insects without any movement were considered dead. Three oil-free flasks were used as control with same number of insects. Three replicates were used for each concentration.

#### **Repellency test**

The abdomen of a pigeon (4×5 cm<sup>2</sup> area) was shaved to expose the skin to the blood starved female mosquitoes. Test oil was applied to skin as pure oil or diluted with ethanol or paraffin oil with the aid of a micropipette at 0.5 and 1 µl/cm<sup>2</sup>; the dose was evenly distributed on the treated area. Controls with and without ethanol or paraffin were used and three replicates were used for each treatment. Treatments were placed against wire netted cages (45× 45×45 cm<sup>3</sup>), each containing more than 250 blood starved female mosquitoes. The time when 2 or more mosquitoes started to bite and feed on the treated area was considered as the protection time, according to Tawatsin *et al.* (2006).

#### **Statistical analysis**

Median lethal (LC<sub>50</sub>) values, 95 % fiducial limits and slopes were estimated according to Finney (1971). Repellency results were analyzed by the Costat 2 statistical program, employing the Student- Newman-Keuls Test to compare among means of treatments.

## **RESULTS AND DISCUSSION**

### **The Chemical composition of the tested oils**

#### **Adulticidal trials**

Results of adulticidal activity are shown in Table 2. The two oils showed similar LC<sub>50</sub> values of 0.215 and 0.217 µl/ l, respectively. At higher concentrations, *M. longifolia* oil was more toxic than *L. dentata* oil, where LC<sub>95</sub> values were 0.309 and 0.695 µl/ l, respectively. Essential oil of *Plectranthus incanus* Link showed LC<sub>50</sub> value of 19.6 µl/ l against *Culex fatigans*, the main compounds in this oil were fenchone, piperitone oxide and piperitenone oxide, where the LC<sub>50</sub> value of the commercial citronella oil was 31 µl/ l (Pal *et al.*, 2011); fenchone was one of the main compounds in *L. dentata* oil used in the present study. Yang *et al.* (2005) reported LC<sub>50</sub> value of 254 µl/ l for *Mentha piperita* oil against *Culex pipiens quinquefasciatus* adults. Wintergreen oil showed knockdown activity against *C. pipiens pallens* adults at 12 µL/L and cinnamon oil exhibited fumigation toxicity against *C. pipiens pallens* adults with the LC<sub>50</sub> value of 0.31 µL/L (Wei-Bin *et al.*, 2013)

**Table 1: Chemical Composition of tested essential oils**

Constituent	R <sub>t</sub>	<i>L. dentata</i>	<i>M. longifolia</i>
(-)-Borneol	27.03		1.72
(+)-Fenchol	22.55	1.74	
(+)-Sabinene	11.70		0.42
1,8-Cineole	16.20		7.35
Camphor	27.55	60.39	
Carvacrol	37.20		
Carvacrol methyl ether	31.78		
Caryophyllene Oxide	53.80		
Cis-Isopulegone	29.51		1.47
dl-Limonene	14.94	0.41	0.30
Eucarvone	42.73		2.68
Fenchone	22.34	28.05	
Gamma-Terpinene	13.44		0.55
Germacrene	45.39		0.20
Linalool	21.72	1.80	
Linalool oxide	19.23	0.47	
L-Menthone	26.87		6.62
Myrtenal	31.16	1.04	
Myrtenol	29.30	0.62	
NI	25.48	1.19	
NI	28.10	0.74	
NI	51.21		0.13
NI	57.12		0.52
NI	81.25		0.23
Para-cymene	16.06		
Para-Cymene-8-ol	30.30	0.27	
p-Cymenene	21.49		
Piperitone	35.53		0.15
Pulegone	34.69		74.95
Trans-caryophyllene	40.53		0.61
Trans-linalool oxide	20.88	0.49	
trans-p-Menthan-3-one	27.68		0.63
trans-β-Ocimene	15.81		
Verbenone	32.78	0.30	
α-Caryophyllene	43.37		0.17
α-Pinene	8.38		0.19
α-terpinene	47.03	0.37	
α-terpinolene	19.09		
β-Bisabolene	46.21		
β-Myrcene	12.74		
β-Pinene	11.8	0.89	0.13
β-Selinene	45.64	0.36	

R<sub>t</sub> = retention time**Table 2: Probit analysis for adulticidal activity of test oils against *C pipiens* females**

Oil	Oils Toxicity		
	LC <sub>50</sub> <sup>a</sup> (95% FL)	LC <sub>95</sub> (95% FL)	Slope ± SE
<i>M. longifolia</i>	0.215 (0.204- 0.225)	0.309 (0.292-0.334)	10.44 ± 0.97
<i>L. dentata</i>	0.217 (0.193- 0.239)	0.695 (0.565- 0.95)	3.25 ± 0.34

LC<sub>50</sub> = μl/L

**Repellency results**

The results of repellent activity of *L. dentata* and *M. longifolia* oils are presented in Table 3. The longest protection time against mosquito biting (165 min) was achieved by *L. dentata* oil at 1 µl/cm<sup>2</sup> when the oil was applied in paraffin oil, while the same dose applied as pure oil or dissolved in ethanol gave 85 min protection time, that is, paraffin oil prolonged the protection time by two fold and there was a significant difference between the two treatments. *M. longifolia* oil, at 1 µl/cm<sup>2</sup>, applied in paraffin oil, gave protection against mosquito attack for only 64.67 min, which means that *L. dentata* oil, at the same dose, is 3 times more effective than *M. longifolia* oil. Moreover, *L. dentata* oil, at 1 µl/cm<sup>2</sup>, without paraffin oil, was more effective than *M. longifolia* oil dissolved in paraffin oil at the same dose, and gave protection against mosquito biting for 85 min, while *M. longifolia* oil protection time was 65 min only with a significant difference between the two treatments. The protection time of the two oils, dissolved in paraffin oil, at 0.5 µl /cm<sup>2</sup>, decreased to be 44.67 and 36 min, respectively. *L. dentata* oil was significantly more effective than *M. longifolia* oil. At the same dose, without paraffin oil, the protection time for the two oils declined to 32.67 and 12 min, respectively. *Zingiber officinals* volatile oil, at 4 mg/cm<sup>2</sup>, showed repellent activity for 120 min against adult *Culex quinquefasciatus* (Pushpanathan, *et al.*, 2008). At 0.005 mg/cm<sup>2</sup>, essential oils of clove and juniper berry gave good repellency against female *Culex pipiens pallens* (Kang *et al.*, 2009). *Lavandula gibsoni* essential oil, at 2 mg/cm<sup>2</sup>, gave 100% protection against *Aedes aegypti* adults for more than 7 h (Kulkarni *et al.*, 2013). Kumar *et al.* (2011) reported that the protection time of *Mentha pipertia* essential oil against bits of *Aedes aegypti* was 150 min when the oil was applied at 4 µl /cm<sup>2</sup>. Peng *et al.* (2009) reported that essential oil from *Zanthoxylum beecheyamum* var. *alatum* provided protection for 6.54 h against *Culex pipens quinquefasciatus* at 1.5 mg/cm<sup>2</sup>. *Cymbopogon citratus* oil gave protection for 84 min against adults of *Culex uinquefasciatus* at 0.21 mg/cm<sup>2</sup> (Phasomkusolsil *et al.*, (2011).

**Table 3: Protection time (min) of test oils against bits of *C pipiens* adults**

Treatment	Protection time (min)
	Mean ± SD*
<i>L. dentata</i> oil in paraffin (1µl/cm <sup>2</sup> )	165 ± 5 <sup>a</sup>
<i>L. dentata</i> oil (1µl/cm <sup>2</sup> )	85 ± 5 <sup>b</sup>
<i>M. longifolia</i> oil in paraffin (1µl/cm <sup>2</sup> )	64.67 ± 5 <sup>c</sup>
<i>L. dentata</i> oil in paraffin (0.5µl/cm <sup>2</sup> )	44.67 ± 1.6 <sup>d</sup>
<i>M. longifolia</i> oil (1µl/cm <sup>2</sup> )	43.67 ± 1.2 <sup>d</sup>
<i>longifolia</i> oil in paraffin (0.5µl/cm <sup>2</sup> )	36 ± 2.65 <sup>e</sup>
<i>L. dentata</i> oil (0.5µl/cm <sup>2</sup> )	32.67 ± 2.5 <sup>e</sup>
<i>M. longifolia</i> oil (0.5µl/cm <sup>2</sup> )	12 ± 2 <sup>f</sup>
Controls	1 ± 0.0 <sup>g</sup>

\*Means with the same letters are not significantly different at the 0.05 level.

## CONCLUSION

The results of the present study showed that the essential oils of tested plant species have insecticidal and repellent activity against adults of *C pipiens*. The oil of *L. dentata*, diluted with paraffin oil, could be used as protection agent against biting of *C pipiens* adults to protect people against disease transmission. Essential oils have many advantages over conventional pesticides; they have pleasant smell, they are blend of many compounds which render them difficult for insect to develop resistance, and they do not cause residue problems in the environment because they are volatile. Further studies are needed to isolate the most effective compounds from these oils. The results indicated that plants are potential safe alternatives to hazardous conventional pesticides, especially for the control of the public health pests.

### Acknowledgement

This work was supported by King Saud University, Deanship of Scientific Research, College of Food and Agriculture Sciences Research Center.

## REFERENCES

- Al-Harby, H. A; Al-Rajhi, D and Hussein, HI. 2008. Pesticidal properties of three Saudi plants: J. Agric. Sci. Mansoura Univ. 33 (10): 7487-7492.
- Ansari, M. A., Vasudevan, P., Tandon, M. and Razadan, R. K., 2000. Larvicidal and mosquito repellent action of peppermint (*Mentha piperita*) oil. Bioresource Technology. 71, 267 – 271.
- Al-Sarar, A. S.; Hussein, H. I; Abobakr; Y; Bayoumi, A. E. and Al-Otaibi, M. T. 2014. Fumigant Toxicity and Antiacetylcholinesterase Activity of Saudi *Mentha longifolia* and *Lavandula dentata* species against *Callosobruchus maculatus*. Turkish Journal of Entomology, 38 (1): Accepted
- Baka, ZAM. 2010. Antifungal activity of six Saudi medicinal plant extracts against five phytopathogenic fungi. Archives of Phytopathology and Plant Protection, 43 :736-743.
- Bousmah, L, Boti, J.B., Bekkara, F.A., Casanova, V.J, 2006. Intraspecific chemical variability of the essential oil of *Lavandula dentata* L. from Algeria. Flavour and fragrance Journal 21, 368 – 372.
- Briassoulis, G., 2001. Toxic encephalopathy associated with use of DEET insect repellents: a case analysis of its toxicity in children. Hum. Exp. Toxicol. 20, 8 – 14.
- Cetin, H., Erler, F. and Yanikoglu, A. 2004. Larvicidal activity of a botanical natural product, AkseBio2, against *Culex pipiens*. Fitoterapia. 75, 724 – 728.
- Cetin, H. and Yanikoglu, A. 2006. A study of the larvicidal activity of *Origanum* (Labiatae) species from southwest Turkey. Journal of Vector ecology. 31 (1): 118 – 122.

- Dob, T., Dahmane, D., Tayeb, B., Chelghoum, Ch., 2005. Chemical composition of the essential oil of *Lavandula dentata* L. from Algeria. *International Journal of Aromatherapy* 15, 110–114.
- Dolan, M.; Dietrich, G.; Panella, N.A.; Montenieri, J. and Karchesy, J. J. 2007. Biocidal activity of three wood essential oils against *Ixodes scapularis* (Acari: Ixodidae), *Xenopsylla cheopis* (Siphonaptera: Pulicidae), and *Aedes aegypti* (Diptera: Culicidae). *J. Econ. Entomol.* 100(2): 622-625.
- Finney, D.N., 1971. *Probit Analysis*, 3<sup>rd</sup> Edition. Cambridge Univ. Press, London, 318 pp.
- Gillij, Y. G., Glesiser, R. M. and Zygadlo, J. A., 2008. Mosquito repellent activity of essential oils of aromatic plants growing in Argentina. *Bioresource Technology*. 99, 2507 – 2515.
- Govindarajan, M. ; Sivakumar, R.; Rajeswary, M. and Yogalakshmi, K. 2013. Chemical composition and larvicidal activity of essential oil from *Ocimum basilicum* (L.) against *Culex tritaeniorhynchus*, *Aedes albopictus* and *Anopheles subpictus* (Diptera: Culicidae). *Experimental Parasitology*, 134 (1): 7–11.
- Hafedh, H., Fethi, B.A., Mejdj, S., Emira, N., Amina, B., 2010. Effect of *Mentha longifolia* L. ssp *longifolia* essential oil on the morphology of four pathogenic bacteria visualized by atomic force microscopy. *African Journal of Microbiology* 4, 1122-1127.
- Hussain, A.I., Anwar, F., Nigam, P.S., Ashraf, M., Gilani, A.H., 2010. Seasonal variation in content, chemical composition and antimicrobial and cytotoxic activities of essential oils from four *Mentha* species. *Journal of the Science of Food and Agriculture* 90, 1827-36.
- Hussein, H.I. 2005. Composition of essential oils isolated from three plant species and their molluscicidal activity against *Theba pisana* snails. *J. Pest Cont. & Environ. Sci.*, 13 (2): 15-24.
- Imelouane, B., Elbachiri, A., Ankit, M., Benzeid, H., Khedid, K., 2009. Physico-Chemical Compositions and Antimicrobial Activity of Essential Oil of Eastern Moroccan *Lavandula dentata*. *International Journal of Agriculture and Biology* 11,113-118.
- Imelouane, B., Elbachiri, A., Wathélet, J.P., Dubois, J., Amhamdi, H., 2010. Chemical composition, cytotoxic and antioxidant activity of the essential oil of *Lavandula dentata*. *World Journal of Chemistry* 5, 103-110.
- Kang, S; Kim, M.; Seo D; Noh,D; Yang, J; Yoon,C. and Kim,G. 2009. Comparative repellency of essential oils against *Culex pipiens pallens* (Diptera: Culicidae). *Journal of the Korean Society for Applied Biological Chemistry*, 52: 353-359.
- Koliopoulos, G., Pitarokili, D., Kioulos, E., Michaelakis, A., Tzakou, O., 2010. Chemical composition and larvicidal evaluation of *Mentha*, *Salvia*, and *Melissa* essential oils against the West Nile virus mosquito *Culex pipiens*. *Parasitology Research* 107, 327–335.

- Kulkarni, R.; Pawar, P. V.; Joseph, M. P. ; Akulwad, A. K. ; Sen, A. and Joshi, S. P. .2013. *Lavandula gibsoni* and *Plectranthus mollis* essential oils: chemical analysis and insect control activities against *Aedes aegypti*, *Anopheles stephensi* and *Culex quinquefasciatus*, Journal of Pest Science, 86: 713-718.
- Kumar S.; Wahab, N and Warikoo, R. 2011. Bioefficacy of *Mentha piperita* essential oil against dengue fever mosquito *Aedes aegypti* L. Asian Pac J Trop Biomed. ;1:85-88.
- Manimaran, A. ; Cruz, M. M. J. J.; Muthu, C; Vincent, S. and Ignacimuthu, S. 2012. Larvicidal and knockdown effects of some essential oils against *Culex quinquefasciatus* Say, *Aedes aegypti* (L.) and *Anopheles stephensi* (Liston). Advances in Bioscience and Biotechnology, 2012, 3: 855-862.
- Michaelakis, A., Theotokatos, S. A., Koliopouls, G. and Chorianoopoulos, N. G. 2007. Essential oils of *Satureja* species: insecticidal effect on *Culex pipiens* Larvae (Diptera: Culicidae). Molecules. 12, 2567 – 2578.
- Morales, M. A., Barrandeguy, M., Fabbri, C., Garcia, J. B., Vissani, A., Trono, K., Gutierrez, G., Pigretti, S., Menchaca, H., Garrido, N., Taylor, N., Fernandez, F., Levin, S., Enria, D., 2006. West Nile Virus isolation from equines in Argentina, 2006. Emerg. Infect. Dis. 12, 1559 – 1561.
- Odeyemi, O.O., Masika, P., Afolayan, A.J., 2008. Insecticidal activities of essential oil from the leaves of *Mentha longifolia* L. subsp. *capensis* against *Sitophilus zeamais* (Motschulsky) (Coleoptera: Curculionidae). African Entomology 16, 220-225.
- Pal, M; Kumar, A; and Tewari, S. K. 2011. Chemical composition and mosquito repellent activity of the essential oil of *Plectranthus incanus* Link. Facta Universitatis ( Physics, Chemistry and Technology), 9: 57 – 64.
- Peng, Z., Beckett, A. N., Engler, R. J. Hoffman, D. R., Ott, N. L., Simons, F.E.R., 2004. Immune responses to mosquito saliva in 14 individuals with acute systemic allergic reactions to mosquito bites. J. Allergy Clin. Immunol. 114, 1189 – 1194.
- Peng YH1, Zhang Y, Zeng DQ, Chen FF, Zhong HY, Li ZH and Huang Y. 2009. Bioactivity and chemical composition of essential oil from *Zanthoxylum beecheyanum* var. *alatum* leaves against *Culex pipiens quinquefasciatus* (Diptera: Culicidae). Ying Yong Sheng Tai Xue Bao. ,20:1488-94.
- Phasomkusolsil, S. and Soonwera, M. 2011. Comparative mosquito repellency of essential oils against *Aedes aegypti* (Linn.), *Anopheles dirus* (Peyton and Harrison) and *Culex quinquefasciatus* (Say). Asian Pacific Journal of Tropical Biomedicine, S113-S118.
- Prajapati, V., Tripathi, A. K., Aggarawal, K. K. and Khanuja, S. P. S., 2005. Insecticidal, repellent and oviposition-deterrent activity of selected essential oils against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus*. Bioresource Technology. 96, 1749 – 1757.



- Pushpanathan .T.; Jebanesan ,A. and Govindarajan , M. 2008. The essential oil of *Zingiber officinalis* Linn (Zingiberaceae) as a mosquito larvicidal and repellent agent against the filarial vector *Culex quinquefasciatus* Say (Diptera: Culicidae). Parasitol Res, 102:1289–1291
- Sardelis M. R., Turell M. J. and Andre R. G. (2003) Experimental transmission of st Louis encephalitis virus by *Ochlerotatus japonicas*. *Journal of the American Mosquito control Association* 19, 159 – 162.
- Service, M. 2004. Medical entomology for students, 3<sup>rd</sup> ed. Cambridge university press, Cambridge, p. 285.
- Smith G. C., Moore C. G., Davis T., Savage H. M., Thapa A. B. Shrestha S. L., and Karabatsos N. (1993) Arbovirus surveillance in northern Colorado, 1987 and 1991. *Journal of Medical Entomology* 30, 257 – 261.
- Sritabutra, D and Soonwera, M. 20013. Repellent activity of herbal essential oils against *Aedes aegypti* (Linn.) and *Culex quinquefasciatus* (Say.). *Asian Pacific Journal of Tropical Disease*, 3: 271-276.
- Tawatsin, A.; Asavadachanukorn,P.; Thavara,U.; Wongsinkongman, P.; Bansidhi, J.; Boonruad, T.; Chavalittumrong,P.;Soonthornchareonnon, N; Komalamisra, N and Mulla , M. S. 2006. Repellency of essential oils extracted from plants in Thailand against four mosquito vectors (Diptera: Culicidae) and oviposition deterrent effects against *Aedes aegypti* (Diptera: Culicidae). *Southeast Asian J Trop Med Public Health*, 37: 915-931.
- Touati, B., Chograni, H., Hassen, I., Boussaid, M., Toumi, L., Ben Brahim, N., 2011. Chemical composition of the leaf and flower essential oils of Tunisian *Lavandula dentata* L. (Lamiaceae). *Chemistry & Biodiversity* 8, 1560-1568.
- Wei-Bin, MA.; Jun-Tao, F.; Zhi-Qing, MA.; Zhi-Li, J. and Xing, Z. 2013. Biological activities of wintergreen oil and cinnamon oil against *Culex pipiens pallens* (Diptera: Culicidae). *Acta Entomologica Sinica*, 56 : 1391-1396.
- WHO. 1992 Vector resistance for pesticides. World health organization technical report serial 818, WHO, Geneva, p. 62
- Yang,P.; Yajun MA and Zheng, S. 2005. Adulticidal Activity of Five Essential Oils against *Culex pipiens quinquefasciatus*. *J. Pestic. Sci.*, 30: 84–89.

**التركيب الكيميائي والنشاط الابادى والطارد للزيوت الأساسية المعزولة من نباتى  
الحبك والخزامى ضد الطور البالغ من بعوض الكيولكس بيبينز.**

على سعيد آل-سرار

قسم وقاية النبات- كلية علوم الأغذية والزراعة- جامعة الملك سعود- ص ب- ٢٤٦٠- الرياض  
١١٤٥١- المملكة العربية السعودية.

تم تقييم النشاط الابادى والطارد للزيوت الأساسية المعزولة من نباتى الحبك والخزامى ضد  
الاناث البالغة من بعوض الكيولكس بيبينز. كان التركيز القاتل لـ ٥٠% من الاناث البالغة هو  
٢١٥ و. ٢١٧. ميكرولتر/ لتر لكلا الزيتين على التوالي. أظهر زيت نبات الخزامى نشاطاً أعلى  
من زيت نبات الحبك كمادة طاردة. أطال زيت البرافين وقت الحماية لكلا الزيتين وكان أطول وقت  
للحماية هو ١٦٥ دقيقة لزيت الخزامى عند تطبيقه بتركيز ١ ميكرولتر / سم مربع فى زيت البرافين.  
كانت المركبات الأساسية فى زيت الحبك هى بوليغون، ١- منثون، ١- و٨- سنيول، يوكارفون،  
بورنيول- بينما كانت المركبات الأساسية فى زيت نبات الخزامى هى كامفور، فنكون، لينالول،  
وفنكول. كان كلا من الزيتين فعالاً كمادة طاردة مقارنة بالزيوت التجارية. هناك حاجة لمزيد من  
الدراسة لعزل أكثر المركبات فعالية من هذه الزيوت.  
الكلمات الدالة: كيولكس بيبينز، حبك، خزامى، مواد طاردة، النشاط الابادى ضد البالغات.