

EFFECT OF CHROMIUM PICOLINATE SUPPLEMENTATION ON GROWTH PERFORMANCE, SERUM BIOCHEMICAL PARAMETERS, AND IMMUNE STATUS OF BROILERS

T. I. Mohamed & G. R. El-Sayed*

Dept. Of Nutrition And Nutritional Deficiency Diseases

*Dept. Of Biochemistry And Chemistry Of Nutrition **

Faculty Of Vet. Medicine, Mansoura University

ABSTRACT

Two hundred and forty broiler chicks (one day old), were randomly divided into four equal groups, each group was subdivided to be in duplicate (30 chicks for each). Four levels of dietary chromium supplementation (0 control, 800, 1600, and 3200 µg /Kg) were provided in the form of chromium picolinate throughout the experimental period (6 weeks). The chicks were fed on starter diet (23% CP & 3200 Kcal ME/Kg) for the first 3 weeks and grower- finisher diet (20% CP & 3200 Kcal ME/Kg) through the next three weeks. Body weight and feed intake were determined and feed conversion ratio was calculated. At the end of experiment, blood samples were collected for obtaining clear sera which were used for determination of the total proteins, albumin, globulin, glucose, triacylglycerol, high density lipoprotein (HDL), low density lipoprotein (LDL), phospholipids and non esterified free fatty acids (NEFFAs). Also, liver function enzymes including aspartate aminotransferase (AST), alanine aminotransferase (ALT) and gamma glutamyl transferase (GGT) were determined. In addition, serum samples were used for determination of haemagglutination inhibition titer against Newcastle disease, γ globulin, IgG and IgM. At end of the experiment, five birds from each replicate were used for determination of carcass traits and composition. The obtained results revealed that feeding high chromium levels (1600 and 3200 µg/Kg) significantly increased ($P<0.05$) the final body weight and average daily weight gain. However, chromium had no effect on feed conversion. The dietary supplements of chromium (1600, 3200 µg/Kg) significantly reduced the blood glucose, the total cholesterol level, LDL and NEFFAs. Also, chromium supplements (1600 and 3200 µg/Kg) increased serum phospholipids, HDL, total proteins, albumin and globulin. The liver function tests (AST, ALT and GGT) indicated no differences between the different groups. High chromium supplementation

(1600 and 3200 µg/Kg) increased the γ-globulin levels. Haemagglutination inhibition antibody titer against Newcastle vaccine. IgG and IgM were increased with increasing the chromium level in the diets. The carcass yield did not affected by chromium supplementation (1600 and 3200 µg/Kg). There were decrease in the abdominal pad and increase in breast yield at the level of 1600 and 3200 µg/Kg of chromium. The protein percent of carcass was increased and fat percent was decreased while the ash and moisture were not significantly affected.

INTRODUCTION

Trivalent chromium is involved in carbohydrate metabolism and recognized as active component of glucose tolerance factor (**Rosebrough and Steele, 1981**). It has been reported that chromium plays an important role in regulating glucose metabolism in human and laboratory animals (**Mertz, 1993**). Chromium facilitates interaction between insulin and insulin receptors (**Mooradian and Morley, 1987**), and involved in lipid, protein and nucleic acid metabolism (**Okada et al., 1984; Ohba et al., 1986; Press et al., 1991; McCarty, 1991; Page, 1991; Lien et al., 1999**). Cell proliferation and synthesis of regulatory protein form the basis of an effective immune response. It can be affected by dietary chromium as improved immune response has been observed when organic forms of chromium were supplemented to stressed feeder calves (**Chang and Mowat, 1992; Moonis-Shageer and Mowat, 1993**) and to dairy cows (**Burton et al., 1993**).

Stresses and diseases both increased urinary excretion of chromium (**Borel et al., 1984; Anderson et al., 1988**) and may exacerbate a marginal chromium deficiency. Therefore, deficiency of trivalent chromium may occur in chicken under stress condition because of metabolic changes mediated by various stressors. Chromium deficiency may slow growth rate, impair glucose tolerance and ultimately lead to diabetes or coronary heart disease (**Simonoff et al., 1984**). So, chromium is recognized as an essential trace element for human and **NRC (1989)** has recommended an intake of 50 to 200 µg /Kg of trivalent chromium for the adult human. However, an appropriate recommendation of the chromium requirement of poultry has not been made (**NRC, 1994**), because of poor absorption of chromium chloride which was largely used as a chromium supplement in the early studies (**Dowling et al., 1989; Lien et al., 1996**). Several investigators have demonstrated that the effect of chromium chloride is insignificant as a dietary supplement (**Li and Stoecker, 1996; Lien et al., 1996**).

Chromium picolinate, an organic and low toxic form of trivalent chromium can stimulate insulin activity (**Evans, 1989; Walker, 1993; Amolkon et al., 1995**). Organic chromium im-

proved immune response and increased the chromium concentration in the kidney and spleen (Anderson and Kozlovsky, 1985; Anderson et al., 1993 ; Mooney and Chromwell 1997). Plant products contain a low content of chromium, implying that broilers may have deficiency of chromium because their diets mostly are of plant ingredients (Gibson 1989). Poultry required higher levels of chromium because β -cells of pancreas of poultry are not being active as those of mammals and secreting a small amount of insulin (Lien et al., 1993; 1996; Ward et al., 1994). Therefore, the present study is to investigate the effects of dietary supplementation levels of chromium picolinate on the growth performance, serum biochemical parameters and immune response.

MATERIALS AND METHODS

Two hundred and forty broiler chicks (one day old), were equally divided at random into four groups, each group was subdivided to be in duplicate (30 chicks for each). Four levels of dietary chromium supplementation (0 control, 800, 1600, and 3200 $\mu\text{g Cr /Kg}$) were provided in the form of chromium picolinate. The broilers were raised for 6 weeks in a pen with natural ventilation. The chicks were fed on starter diet (23% CP & 3200 Kcal ME/Kg) for the first 3 weeks and grower- finisher diet (20% CP & 3200 Kcal ME/Kg) through the next three weeks (table. 1) .

The feed and water were provided ad-libitum. The birds were vaccinated against Newcastle disease virus using Hitchener and LaSota vaccines at 7 and 14 days of age, respectively. Body weight and feed intake were determined and the feed conversion ratio was calculated. At end of the experiment, blood samples were collected from the wing vein of 10 birds from each group (5 birds for each replicate). Blood samples were centrifuged for obtaining clear sera which were used for determination of total proteins (Cornel et al., 1949), albumin (Dumas, 1971), globulin (subtracting albumin from total proteins), glucose (Trinder, 1969), triacylglycerols (Young and Postaner, 1975), total cholesterol (Melattini, 1978), HDL (Clark et al. 1983), LDL (Friedwald et al., 1973), phospholipids (Zilversmit, and Davis, 1950) and NEFFAs (Dnucombe, 1964). Liver function tests including AST, ALT (Reitman and Frankel, 1957) and GGT (Persijn and Vonderk, 1976) were determined. Regarding the immune response, the serum samples were used for determination of haemagglutination inhibition titer against Newcastle disease (Anon, 1980), γ -globulin (Davis and Ornistein, 1964), IgG and IgM (Erhard et al., 1992).

At end of the experiment, five birds from each replicate were weighed, slaughtered and used for studying the carcass traits and composition according to AOAC (1984). The obtained data were analyzed using one way ANOVA test (Senedecor and Cochran, 1989).

Table (1) : Physical and chemical composition of experimental diet

Ingredients	0-3 weeks (%)	3-6 weeks (%)
Ground yellow corn	53.96	63.82
Soybean meal(44%)	30.00	23.10
Fish meal (72%)	7.25	6.12
Soybean oil	5.63	4.00
Limestone	0.91	0.88
Dicalcium phosphate	1.72	1.53
Sodium chloride	0.30	0.30
Vit.& min. premix*	0.25	0.25
DL-Methionine	0.07	.
Calculated value**		
ME (Kcal/Kg)	3200	3200
Crude protein %	23.0	20.0
Calcium %	1.00	0.90
Available phosphate %	0.45	0.40
Lysine %	1.33	1.12
Methionine %	0.50	0.39

* Vitamin and mineral premix per 2.5 Kg supplied the following; Vitamin A, 3.5×10^6 IU; Vitamin D 5×10^5 IU; vitamin E, 25000 IU; vitamin K, 150 mg; riboflavin 1000 mg; pantothenic acid, 3000 mg; niacin, 7000 mg; choline, 3.5×10^5 mg; biotin, 50 mg; folic acid, 150 mg; vitamin B₁₂ 3.0 mg; thiamin, 500 mg; pyridoxine, 800 mg; Cu, 50g, Fe, 100g; Mn, 70 g; Zn, 50 g; iodine, 350 mg and selenium, 150 mg. The producer recommended 2.5 Kg premix/ ton diet).

** Values are calculated according to feed composition table of NRC (1994).

RESULTS AND DISCUSSION

The effects of feeding the diets supplemented with different levels of chromium picolinate on broiler performance are presented in table 2. Feeding the diets supplemented with chromium picolinate at the level of 800 mg had no significance effect on body weight, feed intake and feed conversion, while the high chromium levels (1600 and 3200 μg /Kg) significantly increased ($P<0.05$) the final body weight and the average daily weight gain. **Kim et al., (1996)** reported that supplementation of 100 to 800 mg chromium picolinate /Kg diet did not affect growth performance of broilers. Also, previous investigations found no response in broiler growth performance when dietary supplements of 100 to 400 mg chromium picolinate /Kg were used (**Lein et al., 1993; Ward et al., 1994; Mortozona, et al., 1998; Hegazi et al., 2001**). Furthermore, **Lein et al., (1999)** reported that a dietary level of 800 μg /Kg chromium picolinate did not affect growth performance of broiler but higher levels of chromium improved the body weight and average weight gain. In mammalian studies, dietary supplements of 200 mg chromium picolinate /Kg diet tended to enhance the growth rates of pigs and calves (**Chang and Mowat, 1992; Moonsle-Shageer and Mowat, 1993; Lindemann et al., 1995**). The difference between broilers and mammals in term of effectiveness of chromium supplementation might be attributed to that pancreatic β -cells in poultry not being as active as those in mammals (**Lien et al., 1999**). Increased weight gain could be due to chromium action via reduction of serum cortisol decreasing its catabolic effect (**Brockman, 1986**). In addition to chromium role in metabolic process, it has been reported that chromium affect nuclear protein synthesis (**Okada et al., 1984**). Also, the increased average daily weight gain in broilers fed high chromium-diet may be due to increase daily feed intake, however, chromium did not affect feed conversion ratio (table, 2). Similar results were reported by **Hossain et al., (1998)** who found that organic chromium supplementation increased the body weight gain of broilers but did not affect feed conversion.

The effect of feeding diets supplemented with different levels of chromium on serum biochemical parameters and liver function tests are shown in table, 3. The obtained results exhibited that the high dietary supplements of chromium (1600 and 3200 μg /Kg) reduced the blood glucose level. This may be through stimulation of the biological activity of insulin by increasing insulin-sensitive cell receptors or binding activity (**Anderson et al., 1991; Morris et al., 1993; Ward et al., 1994**). Plasma insulin concentration was reported to be increased linearly, whereas corticosterone concentration was decreased as dietary chromium supplementation increased (**Sahin et al., 2001**). Cortisol promotes gluconeogenesis and reduces glucose utilization (**Weckes, 1991**). Insulin can stimulate glucose uptake and utilization by cells (**Cupo and Donaldson, 1987**). Related studies in human and other animals have indicated that a dietary supplement of chromi-

um reduced the serum glucose (**Lefavi, et al., 1993; Mertz, 1993; Amolkon, et al., 1995; Lein et al., 1996; 1999**).

The results of the present study (table, 3) indicated that chromium supplements (1600 and 3200 $\mu\text{g} / \text{Kg}$) increased serum total proteins, albumin and globulin. These results are supported by **Hegazi et al., (2001)** who found chromium picolinate increased serum total proteins. It has been reported that chromium had an effect on nuclear protein and RNA synthesis (**Okada et al., 1984; Chang and Mowat, 1992**). This increase in protein level may be due to the anabolic effect of insulin.

The dietary chromium picolinate supplements (1600 and 3200 $\mu\text{g} / \text{Kg}$) significantly reduced ($P < 0.05$) the serum levels of total cholesterol, LDL and NEFFAs while increased levels of both phospholipids and HDL. These findings may explain the role of chromium in decreasing the incidence of coronary heart disease in human (**Simonoff et al., 1984**). Similarly, **McCarty (1991)** found that the chromium increased the HDL level of human serum. **Howard et al., (1993)** indicated that insulin increased liver LDL receptors, so reducing the LDL level in serum. **Lein et al., (1999)** noticed that high chromium levels in diets of broilers increased the serum phospholipids while both total cholesterol and triacylglycerol were not affected. On the other hand, **Hegazi et al., (2001)** found that chromium picolinate reduced the serum total lipids and total cholesterol. The chromium acts as activator of the biological activity of insulin which depresses adipocyte lipolysis by reducing the activity of cAMP and hormone sensitive lipase (**Lambert and Jacquemin, 1979**). Moreover, Pigs fed chromium picolinate supplemented ration had lower plasma NEFFAs concentration than control pigs (**Matthews, et al., 2001**). The liver function tests indicated no difference between the different groups in AST, ALT and GGT (table, 3). **Gursoy (2000)** mentioned that organic chromium is more available, safe and has less toxic effect.

The effects of chromium picolinate supplementation on the immune response of broilers are presented in table, 4. High chromium supplementation (1600 and 3200 $\mu\text{g} / \text{Kg}$) increased the γ -globulin levels. Also, haemagglutination inhibition antibody titer against Newcastle vaccine, IgG and IgM were increased with increasing the chromium level in the diet. Similarly, **Chang and Mowat (1992)** found that chromium picolinate supplementation increased total immunoglobulins and IgM. However, **Moonsie-Shageer and Mowat (1993)** reported that chromium supplementation had no effect on IgM and IgG₂ levels but significantly increased IgG₁. Improving the immune response may be due to the effect of chromium on reduction of cortisol, which has immunosuppressive effect and inhibiting production and activities of cytokines and antibodies (**Kelly, 1988; Kegley and Spears, 1995**). In addition, chromium supplementation in stressed animals can prevent urinary losses of zinc, iron, copper, manganese and selenium (**Schrauzer et al., 1986; Anderson et al., 1988**). The deficiencies of such minerals had been re-

ported to lower resistance to diseases (**Bull, 1990**).

Concerning the effect of chromium supplementation on the carcass traits (table, 5), it was found that carcass weight but not carcass yield was increased by chromium supplementation at the levels of 1600 and 3200 $\mu\text{g}/\text{Kg}$ diet. Also, there was a decrease in the abdominal fat pad and an increase in breast yield at the level of 1600 and 3200 $\mu\text{g}/\text{Kg}$ of chromium supplementation. Moreover, the protein percent of carcass was increased and fat percent was decreased while the ash and moisture were not significantly different. **Hossain et al., (1998)** and **Choet (1999)** found that abdominal fat decreased and breast meat yield increased in broilers fed diets supplemented with chromium picolinate at levels of 300 and 500ppb, respectively. **Evans and Bowman (1992)** and **Kim et al., (1996)** demonstrated that the increase in protein content of broiler carcass may be attributed to the effect of insulin on amino acid uptake. Additionally, **Lein et al. (1999)** mentioned that chromium increased the liver lipid accumulation so decreasing accumulation of fat in abdominal pad.

Collectively from the result of the present study, it could be concluded that chromium picolinate supplementation at a level of 1600 ppb for broiler diets has beneficial effects evidenced by improved growth performance and carcass traits as well as improving the immune response. It also has beneficial consequences in terms of enhanced resistance to diseases by enhancing the effectiveness of vaccines.

Table (2) : Effect of chromium picolinate supplementation on performance of broilers (Mean \pm SE)

Items	Supplemental chromium ($\mu\text{g/Kg diet}$)			
	Control (0)	800	1600	3200
Initial body weight	35.04 \pm 0.19	35.18 \pm 0.25	35.1 \pm 0.28	34.8 \pm 0.21
Final body weight (g)	1920 \pm 51.26 ^b	2165 \pm 42.33 ^{ab}	2280 \pm 46.51 ^a	2265 \pm 35.29 ^a
Average daily gain (g)	44.88 \pm 1.29 ^b	50.71 \pm 1.12 ^{ab}	53.45 \pm 0.85 ^a	53.1 \pm 1.65 ^a
Average daily feed intake (g)	86.17 \pm 3.26 ^b	94.83 \pm 3.12 ^{ab}	101.2 \pm 4.28 ^a	98.77 \pm 3.8 ^a
Feed conversion ratio	1.92 \pm 0.25	1.87 \pm 0.19	1.89 \pm 0.28	1.86 \pm 0.25

^{ab} Means in the same row with no common superscript are significantly different ($P < 0.05$).

Table (3) : Effect of chromium picolinate supplementation on serum biochemical parameters (Mean \pm SE) of broiler chickens.

Biochemical parameters	Supplemental chromium ($\mu\text{g/Kg diet}$)			
	Control (0)	800	1600	3200
Glucose (mg/dl)	179.13 \pm 2.1 ^a	166 \pm 2.44 ^a	123.5 \pm 1.78 ^b	124.17 \pm 2.11 ^b
Total proteins (gm/dl)	5.70 \pm 0.22 ^b	6.78 \pm 0.4 ^b	9.97 \pm 0.38 ^a	11.04 \pm 0.17 ^a
Albumin (gm/dl)	2.97 \pm 0.31 ^b	4.15 \pm .35 ^b	6.25 \pm 0.29 ^a	7.33 \pm 0.32 ^a
Globulin (gm/dl)	2.73 \pm 0.28 ^b	2.63 \pm 0.21 ^b	3.72 \pm 0.32 ^a	3.71 \pm 0.25 ^a
Triacylglycerol (mg/dl)	30.51 \pm 1.46	27.97 \pm 1.45	28.4 \pm 1.28	27.68 \pm 1.53
Total cholesterol (mg/dl)	89.66 \pm 2.84 ^a	79.23 \pm 2.31 ^b	76.19 \pm 3.11 ^b	78.35 \pm 1.9 ^b
Phospholipids (mg/dl)	98.69 \pm 2.56 ^b	103.81 \pm 3.1 ^b	119.37 \pm 2.3 ^a	121.23 \pm 2.4 ^a
HDL (mg/dl)	68.83 \pm 1.19 ^{ac}	74.15 \pm 1.36 ^b	82.92 \pm 2.13 ^a	85.15 \pm 1.98 ^a
LDL (mg/dl)	26.16 \pm 0.85 ^a	21.03 \pm 0.72 ^b	19.25 \pm 0.53 ^b	15.09 \pm 0.66 ^c
NEFFA (mg/dl)	26.18 \pm 2.1 ^a	24 \pm 1.60 ^a	14.8 \pm 1.1 ^b	6.6 \pm 0.23 ^c
ALT(U/L) (U/L)	40.21 \pm 2.45	41.33 \pm 1.52	39.33 \pm 1.29	42.09 \pm 1.0 ⁵
AST(U/L)	20.43 \pm 1.80	21.67 \pm 0.69	23 \pm 0.82	23.36 \pm 0.98
GGT (U/L)	8.87 \pm 0.63	9.12 \pm 0.35	9.47 \pm 0.27	9.41 \pm 0.26

^{abc} Means in the same row with no common superscript are significantly different ($P < 0.05$)

Table (4) : Effect of chromium picolinate supplementation on immune response (Mean \pm SE) of broiler chickens.

Items	Supplemental chromium (μg /Kg diet)			
	Control (0)	800	1600	3200
HI titre*	2.93 \pm 0.13 ^c	4.37 \pm 0.23 ^b	7.49 \pm 0.38 ^a	7.41 \pm 0.26 ^a
γ -Globulin (gm/dl)	0.85 \pm 0.07 ^b	0.88 \pm .11 b	1.36 \pm 0.08 ^a	1.44 \pm 0.09 ^a
IgG (mg/dl)	589.14 \pm 25.7 ^c	597.59 \pm 36.7 ^b	845.64 \pm 40.36 ^a	838.58 \pm 17.6 ^a
IgM (mg/dl)	154.21 \pm 8.83 ^c	196.11 \pm 6.79 ^b	238.33 \pm 12.8 ^a	274.18 \pm 17.6 ^a

abc Means in the same raw with no common superscript are significantly different (P<0.05).

* HI values are log2 of the original liters.

Table (5) : Effect of chromium picolinate supplementation the carcass trials of broilers (Mean \pm SE)

Items	Supplemental chromium (μg /Kg diet)			
	Control (0)	800	1600	3200
Carcass weight (gm)	1295 \pm 39.02 ^b	1435 \pm 32.14 ^{ab}	1519 \pm 34.03 ^a	1515 \pm 26.16 ^a
Carcass yield (%)	67.44 \pm 3.29	66.28 \pm 3.35	66.62 \pm 2.94	66.88 \pm 3.68
Abd. Fat pad (% of carcass)	3.84 \pm 0.34 ^a	3.61 \pm 0.28 ^a	2.58 \pm 0.30 ^b	2.62 \pm 0.41 ^b
Breast yield (% of carcass)	20.1 \pm 1.96 ^b	20.7 \pm 1.38 ^{ab}	21.73 \pm 1.64 ^a	21.65 \pm 0.97 ^a
Carcass composition				
Moisture %	63.21 \pm 4.40	62.35 \pm 4.35	64.91 \pm 3.81	63.46 \pm 0.81
Protein %	18.23 \pm 1.74 ^b	18.69 \pm 1.82 ^{ab}	19.89 \pm 1.59 ^a	19.94 \pm 1.63 ^a
Fat %	16.26 \pm 1.30 ^a	16.32 \pm 0.89 ^a	13.66 \pm 1.12 ^b	14.06 \pm 0.92
Ash %	1.8 \pm 0.17	1.79 \pm 0.31	1.82 \pm 0.21	1.89 \pm 0.30

ab Means in the same raw with no common superscript are significantly different (P<0.05).

REFERENCES

- Amoikon, E. K.; Fernandez., J. M.; Southern, L. L.; Thompson, D. L.; Ward, T. L. and Olcott, J. M. (1995)** : Effect of chromium tripicolinate on growth, glucose tolerance, insulin sensitivity, plasma metabolites and growth hormone in pigs. *J. Anim. Sci.* 73: 1123.
- Anderson, R. A.; Bryden, N. A. and Polansky, M. M. (1993)** : Form of chromium affects tissue chromium concentration. *FASEB, J.* 7: A204.
- Anderson, R. A.; Bryden, N. A.; Polansky, M. M. and Deuster, P. A. (1988)** : Exercise effects on chromium excretion of trained and untrained men consuming constant diet. *J. Appl. Physiol.*, 64:249.
- Anderson, R. A. and Kozlovsky, A. S. (1985)** : Chromium intake, absorption and excretion of subjects consuming self-selected diets. *Am. J. Clin. Nut.* 41: 1177- 1183.
- Anderson, R. A.; Polansky, M. M.; Bryden, N. A. and Canary, J. J. (1991)** : Supplemental chromium effects on glucose, insulin, glucagon and urinary chromium losses in subjects consuming controlled low chromium diets. *American J. of Clinical Nutrition*, 54: 909-916.
- Anon (1930)** : Isolation and identification of avian pathogens. American Association of Avian Pathologists, Texas, USA.
- AOAC (1984)** : Official methods of analysis (14th Ed.). Association of official analytical chemists, Washington D.C.
- Borel, J. S.; Majerus, T. C. ; Polansky, M. M.; Mozer, P. B. and anderson, R. A. (1984)** : Chromium intake and Cr excretion of trauma patients. *Biol. Trace Elem. Res.*, 6:317.
- Brockman, R. P. (1986)** : Pancreatic and adrenal hormonal regulation of metabolism. In: L.P. Milligan , W. L. Grovum, and A.D. Dobson (Ed.) *Control of digestion and metabolism in ruminants*. P. 405. Prentice-Hall, Englewood Cliffs, N.J.
- Bull, R. C. (1990)** : Trace element and immunity. *Western Nutr. Conf.* P 33. Calgary, Alberta, Canada.
- Burton, J. L.; Mallard, B. A. and Mowat, D. N. (1993)** : Effect of supplemental chromium on immune responses of Periparturient and early lactation dairy cows. *J. Anim. Sci.* 71; 1532-1539.
- Chang, X. and Mowat, D. N. (1992)** : Supplemental chromium for stressed and growing feeder calves. *J. Anim. Sci.* 70: 559-565.

- Choet, M. (1999)** : Increasing efficiency of lean tissue deposition in broiler chickens. Rural Industry Research & development Corporation Project. NSW. Australia.
- Clark, D. A. ; Kozel, P. R. and Mosser, E. L. (1983)** : Evaluation of a Kit to measure HDL-c in serum. Clin. Chem., 29, 1311.
- Cornel, A. G.; Baradwill, C. S. and David, M. M. (1949)** : Colorimetric determination of serum total protein. Biol. Chem. J.177,751-768.
- Cupo, M. A. and Donaldson, W. E. (1987)** : Chromium and vanadium effect on glucose metabolism and lipids synthesis in the chick. Poultry Sci. 66: 120-126.
- Doumas, B. T. (1971)** : Spectrophotometric Bromocresol blue for determination of albumin in serum. Clin. Chem. Acta, 31, 87-96.
- Dowling, H. J.; Offenbacher, E. G. and Pi-Sunyer, E. X. (1989)** : Absorption of inorganic trivalent chromium from the vascularly perfuse rat small intestine. J. Nutrition. 119: 1138-1145.
- Duncombe, W. G. (1964)** : The colorimetric micro-determination of non-esterified free fatty acids in plasma. Clin. Chem. Acta, 9: 122-125.
- Erhard, M. H.; Quilstrop, I.; Jungling, A.; Kaspers, B.; Schmidt, P. and Kuhman, R. (1992)** : Development of specific enzyme linked immunosorbent antibody assay for detection of immunoglobulins G,M,A. Using monoclonal antibodies. Poultry Sci., 71:302-310.
- Evans, G. W. (1989)** : The effect of chromium picolinate on insulin controlled parameters in humans. International J. of Biophysical and Medical Research, 11:163-180.
- Evans, G. W. and Bowman, T. D. (1992)** : Chromium Picolinate Increases membrane fluidity and rate of insulin internalization. J. of Inorganic chemistry, 46: 243-250.
- Friedwald, W. T. Levy, R. I. and Friedricksonmen, H. (1973)** : Estimation of LDL-c in plasma without ultracentrifugation. Clin. Chem. 18, 499-502.
- Gibson, R. S. (1989)** : Assessment of trace element status in humans. Progress of Food and Nutrition Sci., 13: 67-111.
- Gursoy, U. (2000)** : Chromium in broiler diets. Feed International 21, 24-27. A review.
- Hegazi, S. M.; Magda, Sh. Taha, and Abo-El-Magd, M. K. (2001)** : Influence of supplemented salbutamol and chromium on growth performance and carcass quality of broilers. 2nd International Conference, Faculty of Vet. Medicine Mansoura University.
- Hossain, S. M. Barrento, S. I. and Silva, G. G. (1998)** : Growth performance and carcass utilization

zation of broiler feed supplemental chromium from chromium yeast. *Animal Feed Science and Technology*, 71(3-4): 217-228.

Howard, B. V.; Schneideman, N.; Palkner, B.; Hefner, S. M. and Laws, A. (1993) : Insulin, health behaviours, and lipid metabolism. *Metabolism*, 42 (9, 51): 25-35.

Kegley, E. B. and Spears, J. W. (1995) : Immune response, glucose metabolism, and performance of feeder calves fed inorganic or organic chromium. *J. Anim. Sci.* 73: 2721-2726.

Kelly, K. W. (1988) : Cross-talk between immune and endocrine system. *J. Anim. Sci.* 66: 2095.

Kim, Y. H.; Han, Ibn. K.; Choi, Y. J.; Shin, I. S.; Shae, B. J. and Kang, T. H. (1996) : Effects of dietary level of chromium picolinate on growth performance, carcass quality and serum traits in broiler chicks. *AJAS*, 9: 341-374.

Lambert, B. and Jacquemin, G. (1979) : Inhibition of epinephrine- induced lipolysis in isolated white adipocytes of aging rabbits by increased alpha- adrenergic responsiveness. *J. Lipid Res.* 20: 208-216.

Lefavi, R. G. ; Wilson, J. D.; Keith, R. E.; Anderson, R. A.; Blessing, D. L.; Hames, G. G. and MacMillan, J. L. (1993) : Lipid-lowering effect of chromium (III) nicotinic acid complex in male athletes. *Nutrition Research*, 13,: 239-249.

Lein, T. E.; Chen, S. Y., Chen, C. L. and Yu, C. P. (1993) : The effect of various levels of chromium picolinate on growth performances and serum traits of pigs. *J. of Chinese Society Animal Science*, 22: 349-357.

Lein, T. E.; Chen, S. Y.; Wu, C. P.; Lu, J. J. and Chou, R. G. (1996) : The effect of supplemental chromium picolinate on growth performances, serum traits and carcass characteristics of pigs. *J. of Chinese Society animal science*, 25: 253-260.

Lein, T. F.; Horng, Y. M. and Yang, K. H. (1999) : Performance, serum characteristics, carcass traits and lipid metabolism of broilers as affected by supplement of chromium picolinate. *British Poultry Science*, 40: 357-363.

Li, Y. C. and Stoecker, B. J. (1998) : Chromium and yogurt effects on hepatic lipids and plasma glucose and insulin of lean and obese mice. *Biological Trace element Research*, 9: 233-242.

Lindemann, M. D.; Wood, C. M.; Harper, A. F. Kornegay, E. T. and Anderson, R. A. (1995) : Dietary chromium picolinate additions improve gain: Feed and carcass characteristics in growing finishing pigs and increase litter size in reproducing sows. *J. Anim. Sci.* 73: 457-465.

- Matthews, J. O.; Southern, L. L.; Fernandez, J. M.; Pontif, J. E.; Bidner, T. D. and Odgaard, R. L. (2001)** : Effect of chromium picolinate and chromium propionate on glucose and insulin kinetics of growing barrows and on growth and carcass traits of growing-finishing barrows. *J. Anim. Sci.* 79 (8): 2172-2178.
- McCarty, M. E. (1991)** : The case of supplemental chromium and a survey of clinical studies with chromium picolinate. *J. Applied Nutrition*, 43: 58-66.
- Melattini, F. (1978)** : Colorimetric determination of serum total cholesterol. *Clin. Chem.* 24, 2161- 2165.
- Mertz, (1993)** : Chromium in human nutrition. A review. *J. of Nutrition*, 123: 626-633.
- Mooney, K. W. and Chromwell, G. L. (1997)** : Effect of dietary chromium picolinate supplementation on growth, carcass characteristics, and accretion rates of carcass tissues in growing finishing swine. *J. Anim. Sci.*73:3351-3357.
- Moonsie-Shageer, S. and Mowat, D. N. (1993)** : Effect of level of supplemental chromium picolinate on performance, serum constituents and immune status of stressed feeder calves. *J. Anim. Sci.*, 71: 232-238.
- Mooradian, A. D. and Morley, J. E. (1987)** : Micronutrient status in diabetes mellitus. *J.Clin.Nutr.*45: 877.
- Morris, B. W. ; Gray, T. A.; and MacNeil, S. (1993)** : Glucose- dependent uptake of chromium in human and rat Insulin-sensitive tissues. *Clinical Science*, 84: 477-482.
- Mortozona, Y.; Hatano, K.; Sugawara, N. and Ishibashi, T. (1998)** : Effect of dietary chromium picolinate on growth, carcass quality, and serum lipids of female broilers. *Animal Science and Technology*. 69:7, 659-665.
- National Research Council (1989)** : Recommended dietary allowances. Washington, D.C. National Academy Press.
- National Research Council (1994)** : Nutrition Requirement of Poultry. 9th Ed. Washington, D.C. National Academy Press.
- Ohba, H.; Suzuki, Y. and Ohba, H. (1986)** : Enhancement of ribonucleic acid synthesis by chromium (III)-bound chromatin. *J. of Inorganic Biochemistry*, 27: 113-119.
- Okada, S.; Tuskada, H. and Ohba, H. (1984)** : Enhancement of nucleic RNA synthesis by chromium (III) in regenerating rat liver. *J. of Inorganic Biochemistry*, 21:113-119.
- Page, T. C. (1991)** : Chromium, tryptophan and picolinate in diets for pig and poultry. A dissertation. Louisiana State University, USA.

- Persijn, J. P. and Vander, S. W. (1976)** : Colorimetric determination of γ - glutamyl transaminase. *J.Clin. Chem.Clin. Biochem.*, 14, 421.
- Press, R. I.; Geller, J. and Evans G. and Evans, G. W. (1991)** : The effect of chromium picolinate on serum cholesterol and apolipoprotein fraction in human subjects. *Western J. of Medicine*, 152: 41-45.
- Reitman, S. and Frankel, S., (1957)** : Colorimetric determination of GOT or GPT activity. *Am. J. Clin. Path.* 28,56
- Rosebrough, R. W. and Steele, N. C. (1981)** : Effect of supplemental dietary chromium or nicotinic acid on carbohydrate metabolism during basal starvation and refeeding periods in poultry. *Poultry Science*, 60: 407-417.
- Sahin, K.; Kucuk, and O., Sahin N. (2001)** : Effects of dietary chromium picolinate supplementation on performance and plasma concentrations of insulin and corticosterone in laying hens under low ambient temperature. *J. Phys. Anim. Nutr.* 85, (5-6):142.
- Snedecor, G. W. and Cochran, P. W. (1989)** : *Statistical Methods*, 8th ed., Iowa State University, Press, Ames, I.A.
- Shrauzer, G. N.; Shrestha, K. P., Molenaar, T. B. and Meade, S. (1986)** : Effect of Cr supplementation on food utilization and the trace element composition in the liver and heart of glucose- exposed young mice. *Biol. Trace Element Res*, 9: 79.
- Simonoff, M.; Llabador, Y. G.; Hamon, A.; Peers, M. and Siminoff, C. M. (1984)** : Low plasma chromium in-patient with coronary artery and heart diseases. *Biological Trace Element Research*, 6:431-439.
- Trinder, P. (1969)** : Determination of blood glucose using an oxidase-/74 peroxidase system with a non carcinogenic chromagen. *J.Clin. Path.*22, 158-161.
- Walker, M. (1993)** : Chromium; The essential mineral. *Health of Foods Business* 5: 51-52.
- Ward, T. L.; Southern, L. L. and Boleman, S. L. (1994)** : Effect of dietary picolinate on growth, nitrogen balance and body composition of growing broiler chickens. *J. Anim. Sci.*, 73 (S.1) : 37.
- Weekes, T. E. (1991)** : Hormonal control of glucose metabolism. In: T. Tsuda; Y.Sasaki; and R. Rawashima (ed.) *Physiological aspect of digestion and metabolism in ruminants*. Proceeding in 7th Int. Symposium on Ruminant Physiology. P183.
- Young, D. and Postaner, L. (1975)** : Enzymatic and colorimetric test for determination of triglycerides. *Clin. Chem.* 21, 5.

Zilversmit, D. B. and Davis, A. K. (1950) : Micro-determination of phospholipids by trichloroacetic acid precipitation. *J. Lab. Clin. Med.* 35, 155-160.

الملخص العربى

تأثير إضافة بيكلونات الكروم على النمو والقياسات الكيميائية الحيوية وخصائص الذبيحة والمستوى المناعى فى دجاج التسمين

طارق إبراهيم محمد و جهاد رمضان السيد*

قسم التغذية وأمراض سوء التغذية، وقسم الكيمياء الحيوية وكيمياء التغذية*

كلية الطب البيطرى - جامعة المنصورة

أجريت هذه الدراسة على مائتين وأربعين من كتاكيت التسمين (عمر يوم) قسمت إلى أربعة مجموعات متساوية وقسمت كل مجموعة إلى تحت مجموعتين متساويتين فى كل منها ثلاثين كتكوتاً، وغذيت الكتاكيت على علائق تحتوى على بيكلونات الكروم بنسب (صفر للمجموعة الضابطة ٨٠٠، ١٦٠٠، ٣٢٠٠ ميكروجرام / كجم) وتم تربيتها لمدة ستة أسابيع وعين الوزن النهائى وكمية العلف المستهلك وكذلك معدل التحويل الغذائى. فى نهاية التجربة أخذت عينات من الدم لعمل القياسات البيوكيميائية، وأيضاً تم ذبح عشرة طيور من كل مجموعة لدراسة مواصفات الذبيحة وتحليلها كيميائياً.

وقد أظهرت النتائج أن التغذية على المستويات العالية (١٦٠٠، ٣٢٠٠ ميكروجرام / كجم) من بيكلونات الكروم أدت إلى حدوث زيادة معنوية فى الوزن النهائى وكذلك فى متوسط الزيادة اليومية فى الوزن، ولم يؤثر على معدل التحويل الغذائى.

أما بالنسبة لتأثير الكروم على القياسات البيوكيميائية الحيوية فى مصل الدم فقد أظهرت النتائج أن التغذية على المستويات العالية من الكروم (١٦٠٠، ٣٢٠٠ ميكروجرام / كجم) أدت إلى نقص معنوى فى نسبة الجلوكوز والكلوسترول الكلى والليپوبروتينات منخفضة الكثافة وكذا نسبة الأحماض الدهنية الحرة وعلى النقيض أدت هذه المستويات العالية إلى زيادة نسبة الفسفوليبيدات والبروتينات الكلية والألبومين ونسبة الجلوبيولين، ولم تحدث تغيرات معنوية فى إنزيمات كفاءة الكبد (الأسبرتات ترانس أمينيز والجلوتامات ترانس أمينيز والجاما جلوتاميل ترانسفيريز).

وجد أن تأثير إضافة الكروم على الزيادة فى وزن الذبيحة كان واضحاً حيث أنها زادت بزيادة كمية الكروم المضافة (١٦٠٠، ٣٢٠٠ ميكروجرام / كجم). وحدث نقص معنوى فى نسبة دهون منطقة البطن وزيادة فى عضلات منطقة

الصدر، ووجد أن هناك زيادة فى نسبة البروتينات ونقص فى نسبة الدهون فى الذبيحة ولكن لم يحدث تغير فى نسبة الرطوبة وكذا نسبة الأملاح.

كما وجد أن زيادة الكروم المضاف (١٦٠٠ ، ٣٢٠٠ ميكروجرام / كجم) أدى إلى زيادة نسبة الجاما جلوبيولين، بالإضافة إلى زيادة نسبة إختبار منع التجلط ضد التحصين لمرض النيوكاسل وكذا نسبة IgG and IgM بزيادة نسبة الكروم وحسنت المستوى المناعى للدجاج.

وقد أوضحت النتائج عامة أن التغذية على المستويات العالية (١٦٠٠ . ٣٢٠٠ ميكروجرام / كجم) من بيكلونات الكروم أدت إلى زيادة معنوية فى الوزن النهائى وكذلك فى متوسط الزيادة اليومية فى الوزن وأدت إلى إعتدال مستوى قياسات التمثيل الغذائى والمؤشرات البيوكيميائية كما أدى إلى تحسين المستوى المناعى.