

GENETIC GAINS AND HETEROSIS FOR SOME EGG PRODUCTION TRAITS IN TWO LINES OF NORFA CHICKENS

A.A. Enab, M.E.Soltan, O. A. El-Weshahy and F.H. Abdou

Dept. of Poultry Production, Fac. of Agric., Minufiya Univ.

(Received: May. 23 , 2010)

ABSTRACT: *The ultimate goal of this study was to develop two lines of Norfa hens (i.e. line egg number at 42 weeks of age (EN) and line body weight (BW) at maturity (38WK)] by using a selection independent culling level method during three generations of selection and crossing two purelines to get hybrid vigor for some egg production traits. A control line was randomly formed from the base population before choosing the individuals of selected line.*

Generally the birds of (EN) line excelled those of (BW) line in both of ASM and EN while the birds of (BW) line were obviously heavier and laid the heaviest eggs comparing to birds of (EN) and control lines. The realized genetic gain for egg number traits during three generations of selection in (EN) line ranged from 10.2 to 26 eggs while expected genetic gain for the same trait in the same line ranged from 1.1 to 2.5 eggs. The realized genetic gains for (ASM) during three generations of selection in Norfa chickens in (EN) line ranged from -7.3 to -16.7 d while expected genetic gains for the same trait in the same line ranged from -0.66 to -1.8 d. All F_1 hybrids for studied traits exceeded the mid-parents in this study. All F_1 crosses for studied traits had positive heterosis values except F_1 crosses for age at sexual maturity (ASM) had negative heterosis value in this study. Heterosis percentages in crossline (BW \times EN) for some egg production traits (i.e. ASM, BW_{SM} , BW_M , EW_{SM} , EW_M , EN_{90d} , EN_{42wk} and EN_{52wk}) were -5.1, 2.4, 6.5, 6.1, 3.1, 14.7, 13.4 and 9.2%, while these estimates in crossline (EN \times BW) for the same traits were -2.5, 2.3, 3.9, 5.1, 2.9, 13.4, 6.4 and 3.9%, respectively.

From the previous results the parental lines EN and BW proved to exploited both additive and non additive variations and could be used to produce superior crosses for egg number and body weight.

Key words: *Genetic gains and heterosis in norfa chickens*

INTRODUCTION

In Egypt, a lot of efforts have been done to improve indigenous chickens. The Egyptian indigenous breeds of chickens have many advantageous such as their high adaptability to local environment and genetic resistance to some serious diseases such as Marek beside the highly acceptable taste and favorable flavor for their meat and egg products.

El-Hadad (2003) found that means of age at sexual maturity (ASM) after four generation of selection for selected and control lines of Norfa chickens were 166.8 and 174.2 d, while these means for egg number till 42 weeks of age (EN_{42wk}) were 68.5 and 63.6 eggs, respectively. Abou El-Ghar and Abdou (2004) found that the egg numbers in the first ninety days of laying (EN_{90d}) were 55.4 and 48.1 eggs for two selected lines of Norfa layers (i.e. egg number and egg weight).

Enab *et al.* (2000) found that the actual and expected genetic gains for EN_{90d} after to generations in Norfa chickens were 8.8 and 2.4 eggs, while these means were 10.8 and 4.7 for EN_{42wk}, respectively. El-Hadad (2003) found that absolute genetic gains for EN_{42wk} in Norfa chickens were 5.8, 4.9, 8.4 and 4.9 eggs in G₁, G₂, G₃ and G₄, respectively.

Many investigators confirmed the superiority of crossbreds over the purebreds regarding egg production traits and some economic traits (Kosba *et al.* 1981; Farghaly and Saleh, 1988; Abdou, 1992; Nawar and Abdou, 1999, and Abou El-Ghar *et al.*, 2007). Crossing between native and foreign breeds had performed better than pure ones (Farhaly and Saleh, 1988, Nawar and Bahie El-Deen, 2000 and Amin, 2008). Abou El-Ghar (2003) showed that both dominance and epistasis were important in heterosis for egg production traits in Norfa strain crosses.

The main purpose of the present study is to improve Norfa strain by using a selection method of independent culling level during three generations of selection to develop two lines of Norfa strain (i.e. egg number "EN" and body weight "BW") to increase the genetic variations. In generation four, crossing the two lines of Norfa strain was done to exploit additive variations in the hybrid vigor in Norfa performance.

MATERIALS AND METHODS

I-Plan and management

The present experiment has been carried out at the Poultry Farm of the Faculty of Agriculture, Minufiya University, Shebin El-Kom, Egypt as a part of Norwegian-Egyptian project "NORFA" for improving hens during four generations through the hatching seasons 1997-2001. In the base population a total of 415 dams of Norfa chickens were divided into three lines based on egg number and body weight at maturity (165 EN, 150 BW and 100 control hens) to produce the next generation, whereas each three dams were mated artificially to a cock. The ultimate goal of this study was to develop two lines of Norfa hens (i.e. EN and BW) by using a selection independent culling level method during three generations of selection and crossing two purelines to get hybrid vigor for egg production traits.

Numbers of hens per generation, line, crosslines and control were shown in Table (1). Artificial insemination was done as a mating system during all generations, and the semen was collected from cocks and inseminated fresh and undiluted into dams. Each sire artificially inseminated three dams in

Genetic gains and heterosis for some egg production traits.....

each generation. Mating of relatives was avoided. Insemination started one week before collecting hatching eggs, each dam was inseminated twice a week. Fertile eggs were collected daily for a couple of weeks and stored in a prepared refrigerator.

Cockerels were separated from pullets in brooding house at 8 weeks of age and at 14 weeks, cockerels moved to individual cages in cock's house while, pullets were moved to individual cages in laying house at 16 weeks of age.

Pullets were fed a starter ration contained 18% crude protein and 2833 kcal. ME/kg. ration until 18 weeks of age, from 19 weeks of age to the end of production period, a layer ration contained 16.5% crude protein and 2758 kcal. / kg. ration.

II. Studied traits.

The following traits were studied:

1. Body weight at sexual maturity (BW_{SM}), in grams.
2. Body weight at 38 weeks of age (BW_M), in grams.
3. Age at sexual maturity (ASM): number of days at the first egg laid.
4. Egg number (EN)
 - 4.1. Number of eggs in the first ninety days of laying (EN_{90d}).
 - 4.2. Number of eggs at 42 weeks of age (EN_{42WK}).
 - 4.3. Number of eggs at 52 weeks of age (EN_{52WK}).
5. Egg weight (EW)
 - 5.1. The average weight of 5 eggs at sexual maturity (EN_{SM}), in grams.
 - 5.2. The average weight of 5 eggs during 35-38 weeks of age (EW_M), in grams.

III- Selection procedures:

During three generations of selection the independent culling levels procedure was applied in the base population to divide it into three lines (i.e. EN, BW and control). The egg number line (EN) was determined by using the overall mean for egg number at 42 weeks of age of the base population plus one standard deviation ($\mu+1$ S.D.), while the body weight line (BW) was determined by using the overall mean of body weight at maturity (38 WK) of the base population plus one standard deviation ($\mu+1$ S.D.). All individuals that failed to come down of these two levels were discarded. Moreover, those selected individual hens should attain at least the general averages of the base population concerning the other studied traits. The same selection method was applied in each of the following generations. Also, the cocks of the base population were divided into these three lines upon this procedure. The birds of the control line were chosen randomly before applying the independent culling levels in the base population to choose the individuals of the two selected lines (i.e. EN and BW).

In the fourth generation two crosslines (i.e. EN x BW and BW x EN) were obtained by crossing EN and BW lines.

Table 1

Genetic gains and heterosis for some egg production traits.....

IV- Statistical analysis:

The statistical analysis was performed using linear models procedure of the statistical analysis system computer program (SAS, 1994). Duncan's new multiple range test was used to compare every two means of different traits studied (Steel and Torrie, 1960).

1. The statistical model:

The statistical model used for analyzing egg production traits between lines in different generation was as follows.

$$Y_{ijk} = \mu + L_i + G_j + LG_{ij} + e_{ijk}.$$

Where:

Y_{ijkl} = Observation on the ijk^{th} chick.

μ = Overall mean,

L_i = Effect of i^{th} line.

G_j = Effect of j^{th} generation.

LG_{ij} = Interaction effect between i^{th} line and j^{th} generation.

e_{ijk} = Random error component, assumed to be normally distributed.

2. Genetic gains and heterosis:

Realized genetic gains (ΔGR) were calculated as a deviation of the mean of parental line from the control mean.

$$\Delta GR = \bar{x}_s - \bar{x}_c$$

Where:

\bar{x}_s = Mean of parental line,

\bar{x}_c = Mean of control.

The expected genetic gains (ΔGE) were calculated according to the formula given by (Prichener's, 1979) as follow.

$$\Delta GE = i \cdot h_j \cdot \delta_A$$

Where:

i = Selection intensity.

h_j = Square root of heritability for the j^{th} trait.

δ_A = Genetic standard deviation of the j^{th} trait.

Average degree of heterosis (ADH %) based on the mid-parents (MP) was determined according to equation given by (Sinha and Khanna, 1975) as follows.

$$ADH\% = \bar{F}_1 - MP/MP \cdot 100$$

Where:

\bar{F}_1 = Mean of crossbred,

MP = mid-parents.

RESULTS AND DISCUSSION

1- Means

Means of lines and crosslines for egg production traits (ASM, BW_{SM} , BW_M , EW_{SM} , EW_M , EN_{90d} , EN_{42wk} and EN_{52wk}) are presented in Table (1). The averages of ASM during generation one were 147.8, 162.3 and 155.1d for EN, BW and control lines while the corresponding averages during generation two were 143.5, 169.2 and 157.1d, respectively. The results shown during the

first and second generations were in a good agreement with those reported by Enab (2001), Abou-Elwewa (2004), Abou El-Ghar and Abdou (2004) and Ben Naser (2007). Whereas they recorded that averages of (ASM) for Norfa pullets ranged from 150.6 to 176.5 d. Means of BW_{SM} during generation one were 1118.8, 1369.4 and 1095.6g for EN, BW and control lines, while the corresponding averages during generation two were 1067.5, 1457.1 and 1110.3g, respectively. Many researchers reported that the average of Body weight at sexual maturity (BW_{SM}) in Norfa chickens ranged from 1005.0 to 1459.0 g (Enab *et al.*, 2000, El-Hadad, 2003, Abou El-Ghar, 2003, Abou-Elwewa, 2004 and Ben Naser, 2007).

The data in Table (1) also show that means of EW_{SM} during generation one were 36.5, 39.6 and 35.4g for EN, BW and control lines, while the corresponding averages during generation two were 35.7, 42.4 and 36.5g, respectively. Similar results were reported by (El-Hadad, 2003; Abou El-Ghar and Abdou, 2004, Ben Naser, 2007 and Abou-Elwewa, 2010) who showed that the average of EW_{SM} in Norfa chickens ranged from 35.6 to 39.1g. The averages of EN_{90d} during the generation one were 60.1, 52.2 and 49.9 eggs for EN, BW and control lines while the corresponding averages during the generation two were 65.3, 50.3 and 48.7 eggs, respectively. Similar results were reported by (El-Hadad, 2003; Abou-Elwewa, 2004 and Abou El-Ghar and Abdou, 2004) who showed that the average of EN_{90d} in Norfa chickens ranged from 49.3 to 64.8 eggs.

There were highly significant differences between lines, generations and lines \times generations interaction for all studied egg production traits (Table, 2).

2- Crossing

In generation three, means of age at sexual maturity (ASM) for two parental lines of Norfa strain (i.e. EN and BW) and control are given in Table (1). It was noticed that the (EN) parental lines had the lowest sexual maturity means (136.9d), while these means were 173.3 and 153.6 d for (BW) parental line and control, respectively.

In generation four, means of (ASM) for two crosslines for Norfa strain (i.e. EN and BW) and control are shown in Table (1). It was clear that the crossline (BW \times EN) had the lowest sexual maturity mean (147.2), while these means were 151.2 and 156.9 for (EN \times BW) cross line and control, respectively. The results in Table (1) show that (EN) parental line had the lowest sexual maturity means (136.9 d) than (BW) parental line and F_1 crosses. Generally, means of ASM in F_1 crosses were less than the parental means (149.2 vs. 155.1d). However, most crossbreds were earliest in reaching sexual maturity compared to pure breeds. Similar results were reported by Nawar (1995), El-Salamony (1996), Nawar and Abdou (1999) and El-Tahawy (2000).

In generation three, averages of body weight at maturity were 1081.8, 1724.5 and 1195.3g for EN, BW parental lines and control, respectively, (Table 1). In generation four, these means in the F_1 crosses were 1457.6, 1494.5 and

Genetic gains and heterosis for some egg production traits.....

Table 2

1230.6g for (EN × BW), (BW × EN) crosslines and control, respectively. However, the (BW) parental line had significantly heavier body weight (1724.5g) than (EN) parental line and F₁ crosses. Generally, means of body weight at maturity in F₁ crosses were heavier than the parental means (1476.1 vs. 1403.2g) (Table, 1).

In generation three, the data listed in Table (1) show that the means of egg weight at maturity (EW_M) were 47.6, 56.2 and 46.1g for EN, BW parental lines and control, respectively. In generation four, the means of egg weight at maturity (EW_M) in the F₁ crosses were 53.4, 53.5 and 47.9g for (EN × BW), (BW × EN) crosslines and control, respectively. It was noticed that, (BW) of parental line had significantly heavier body weight (56.2g) than (EN) parental line and F₁ crosses. Generally, means of body weight at maturity in F₁ crosses were higher than the parental means (53.45 vs. 51.9) (Table, 1).

Generally, crosslines gave heavier body and egg weights at maturity than the native purelines. These results were in agreement with those reported by Nawar (1995), El-Salamony (1996), Nawar and Abdou (1999) and Abou El-Ghar (2003).

In generation three, Table (1) shows that the means of egg number at 42 weeks of age (EN_{42wk}) were 96.6, 70.1, 73.9 egg for EN, BW pure parental lines and control, respectively. In generation four, these means in F₁ crosses were 88.8, 94.6 and 72.5 eggs for (EN × BW), (BW × EN) crosslines and control, respectively. However, the (EN) parental line had significantly higher egg number at 42 weeks of age (96.6 eggs) than (BW) parental line and F₁ crosses. Generally, averages of (EN_{42wk}) in F₁ crosses were higher than the parental means (91.7 vs. 83.4 eggs) (Table, 1). In this respect Kosba *et al.* (1981), Wang and Prichner (1991), El-Hossari and Dorgham (1992) and Nawar and Abdou (1999) reported that, the strain crossing increased rate of laying.

3- Genetic gains

Table (3) presents the realized and expected genetic gains for egg production traits in EN and BW lines of Norfa strain during generations 1, 2 and 3. In EN line, the results in Table (3) show that the realized genetic gain for (ASM) in generations one, two and three were -7.3, -13.6 and -16.7d, while the expected genetic gains were -1.8, -1.2 and -0.66, respectively. In BW line, the results show that the realized genetic gains for (ASM) during the generations one, two and three were 7.2, 12.1 and 19.5d, while the expected genetic gain were 1.4, 0.99 and 0.47, respectively. These results showed that EN line reached sexual maturity earlier than the BW line. This is a result of indirect response for selecting to egg number in this line. These results agree with those obtained by Enab *et al.* (2001) who found that actual genetic gain for sexual maturity in different lines of W. Leghorn ranged from -2.88 to 2.95d, while expected genetic gain ranged from -3.99 to -2.30d. Abou El-Ghar (2003) reported that the realized genetic gains for age at sexual maturity in four lines (SM, EN, BW and EW) of Norfa strain were -26.0, -19.0, -15.0 and -11.0d, while the expected genetic gains for the same traits in the same lines were -0.10, -0.07, -0.06 and -0.05d, respectively.

Genetic gains and heterosis for some egg production traits.....

Table 3

Table 3

Genetic gains and heterosis for some egg production traits.....

In EN line, the results in Table (3) show that the realized genetic gain for (BW_{SM}) during the generations one, two and three were 23.2, -24.8 and -50.3g, while the expected were 13.6, -11.9 and -8.7g, respectively. In BW line, the realized genetic gain for (BW_{SM}) in the first, second and third generations were 273.8, 346.8 and 416.0g, while the expected were 17.4, 14.8 and 8.6g, respectively. These results agreed with those obtained by El-Sakka (1999), Enab *et al.* (2000) and Abou El-Ghar (2003) who reported that the realized genetic gains in (BW_{SM}) ranged from -80.8 to 367.0, while the expected genetic gains ranged from -0.6 to 0.23 in different lines of Norfa chickens.

In EN line, the results in Table (3) show the realized genetic gains for (EW_{SM}) in the generations one, two and three were 1.1, -0.8 and 0.2g, while the expected genetic gains were 0.91, -0.64 and 0.31g, respectively.

In BW line, the results in Table (3) show the realized genetic gains for (EW_{SM}) in the first, second and third generations were 4.2, 5.9 and 9.3g, respectively. Moreover, the expected genetic gains for (EW_{SM}) were 1.2, 0.74 and 0.35g in G_1 , G_2 and G_3 , respectively. These results agreed with those obtained by Abdou *et al.* (1997) and Abou El-Ghar (2003) who reported that the realized genetic gains in (EW_{SM}) ranged from -2.0 to 2.8g, while the expected genetic gains ranged from -0.32 to 1.12g in different lines of Norfa chickens.

In EN line, the results in Table (3) show the realized genetic gains for (EN_{90d}) in generations one, two and three were 10.2, 16.6 and 19.6 eggs, respectively, while the expected genetic gains for the same trait of the same line were 2.5, 2.4 and 1.2 eggs, in the same order.

In BW line, the results in Table (3) showed the realized genetic gains for (EN_{90d}) during the generations one, two and three were 2.3, 1.6 and -1.4 eggs, while the expected genetic gains for the same trait in the same line were 2.1, 2.0 and -1.1 eggs, respectively. The same results were obtained by Sherif (1991), El-Wardany *et al.* (1992), Enab *et al.* (2000) and Abou El-Ghar (2003) who reported that the realized genetic gains in (EN_{90d}) ranged from 3.2 to 11.0 eggs, while the expected genetic gains ranged from 0.02 to 2.4 eggs in different lines of Norfa chickens.

4- Heterosis

Heterosis percentages of some egg production traits are presented in Table (4).

The expressions of averages degree of heterosis (ADH %) based on mid-parent for age at sexual maturity (ASM) in generation four are presented in Table (4). F_1 hybrids were earlier than the mid-parents in reaching sexual maturity. Therefore, negative heterosis values were obtained for these hybrids and the estimated (ADH %) were -2.5 and -5.1 % for (EN × BW) and (BW × EN) crosslines, respectively. The results of heterosis percentages for age at sexual maturity showed negative heterosis effects for all crosses (EN × BW) and (BW × EN). Similar results were obtained by Nawar (1995), Bordas *et al.* (1996), El-Salamony (1996) and Nawar and Bahie El-Deen (2000).

Table 4

Genetic gains and heterosis for some egg production traits.....

In generation four, results of heterosis (ADH %) for body weight at sexual maturity are presented in Table (4). It showed that means of F₁ crosses had positive heterosis values. The estimates of heterosis for crosses EN × BW and BW × EN were 2.3 and 2.4 %, respectively. Similar results were obtained by Zatter (1994) and El-Salamony (1996).

In generation four, average degrees of heterosis (ADH %) for (EW_{SM}) are presented in Table (4). It noticed that egg weight at sexual maturity (EW_{SM}) showed positive heterosis effects in the crosses EN × BW (5.1 %) and BW × EN (6.1 %). These results are in agreement with those reported by El-Salamony (1996) and Nawar and Bahie El-Deen (2000).

In generation four, results of heterosis (ADH %) for egg number in the first 90d of laying (EN_{90d}) are presented in Table (4). It showed that means of F₁ crossed had positive heterosis values. Generally, means of (EN_{90d}) in F₁ crosses were higher than the parental means. The estimated of heterosis (ADH %) for crosses EN × BW and BW × EN were 13.4 and 14.7 %, respectively. These results are in agreement with those reported Zatter (1994), Nawar and Bahie El-Deen (2000), Abou El-Ghar *et al.* (2007) and Amin (2008).

In conclusion, the results of the present study indicate that crossbreeding between two lines of Norfa chickens (i.e. developed for egg number; EN and body weight; BW) is considered an effective way to improve some of egg production traits.

REFERENCES

- Abdou, F. H. (1992). Improve Minufiya chicken through developing the new strain (NORFA). Panel Meeting, Shibin El-kom, Minufiya Province, April 28, 1992.
- Abdou, F. H., A. A. Enab and H. El-Ibiary (1997). Application of ten selection indices for genetic improvement of Norfa chickens. PSA 86th Annual meeting, 3-6 August, Univ., Georgia. Vol. 76 Supplements 1 No. (140) pp 139.
- Abou Elewa, E. (2004). Selection for general immune response and its relation to some economic traits in chickens. M. Sc. Thesis, Fac. Agric. Minufiya Univ., Egypt.
- Abou Elewa, E. (2010). Some genetic parameters of the immune response trait and its utilization in different selection methods in chickens. Ph. D. Thesis, Fac. Agric. Minufiya Univ., Egypt.
- Abou El-Ghar, R. (2003). Combining ability and genetic gain of some economic traits in Norfa chickens. Ph. D. Thesis, Fac. Agric. Minufiya Univ., Egypt.
- Abou El-Ghar, R. and F. H. Abdou (2004). Evaluation of genetic variance components based on the concept of generation means for some economic traits in chickens. Egypt. Poult. Sci., 24: 687 – 704.

- Abou El-Ghar, R.; Hedaia M. Shalan and Hanan H. Ghanem (2007). Matrouh as a common parent in crossing with some local strains of chickens. *Egypt. Poultry Sci. J.*, 27: 805 – 815.
- Amin, E. (2008). Effect of crossing between native and a commercial chicken strain on egg production traits. *Egypt. Poultry Sci. J.*, 28: 327 – 349
- Ben Naser, K. (2007). Using selection to improve some economic traits in Norfa chickens. Ph. D. Thesis, Fac. Agric. Minufiya Univ., Egypt.
- Bordas, A., P. Merat and F. Minvielle (1996). Heterosis in egg laying under divergent selection for residual feed consumption. *Poult. Sci.*, 75: 20-24.
- El-Hadad, E. (2003). Efficiency of selection in laying hens by using supplementary information on feed consumption. Ph. D. Thesis, Fac. Agric. Minufiya Univ., Egypt.
- El-Hossari, M. A. and S. A. Dorgham (1992). Economic evaluation of local chicken crosses under control environment-houses. *Egypt. Poultry Sci. J.*, 12: 843-866.
- El-Sakka, M. M. (1999). Expected and realized genetic gains of some egg quality traits in chickens. M. Sc. Thesis, Fac. of Agric. Minufiya Univ., Egypt.
- El-Salamony, A. L. (1996). A study of heterosis in some egg production traits in Norfa layers. M. Sc. Thesis, Fac. of Agric. Minufiya Univ., Egypt.
- El-Tahawy, W. S. A. (2000). Genotypical improvement of some productive and reproductive traits in local chicken. M. Sc. Thesis, Alexandria Univ., Egypt.
- El-Wardany, A., M. Soltan and F. H. Abdou (1992). Relative efficiencies of different selection methods for improving some egg production traits in Norfa chickens. *Minufiya J. Agric. Res.*, 17: 1791 – 1832.
- Enab, A. A. (2001). Genetic progress achieved in residual feed consumption after generations of selection in Norfa layer chickens. *Egypt. Poult. Sci.*, 21(1): 209 – 220.
- Enab, A. A., N. Kolstad and F. H. Abdou (2001). Multi-source multi-trait selection indices for maximizing genetic improvement of laying hens. *Egypt. Poult. Sci. J.*, 21: 277 – 290.
- Enab, A. A., A. M. El-Wardany and F. H. Abdou (2000). Genetic aspects of egg production traits in Norfa layers under different methods of selection. *Egypt. Poult. Sci. J.*, 20: 1017 – 1030.
- Farghaly, M. H. and K. Saleh (1988). The effect of crossbreeding on egg traits in laying hens. *Egypt. Poult. Sci.*, 8 :376 – 392.
- Kosba, M. A., T. H. Mahmoud , A. Z. Kalil and G. M. Abd-Alla (1981). A comparative study of four breeds of chickens and their FI crosses. *Agric. Res. Rev.*, Cairo, 59: 93-103.
- Nawar, M. (1995). A comparative study of some productive traits between some native and foreign breeds of chickens. Ph. D. Thesis, Fac. of Agric., Minufiya Univ., Egypt.

Genetic gains and heterosis for some egg production traits.....

- Nawar, M. and F. H. Abdou (1999). Analysis of heterotic gene action and maternal effect in crossbred Fayoumi chickens. *Egypt. Poult. Sci.*, 19: 671 – 689.
- Nawar,, M. E. and M. Bahie El-Deen (2000). A comparative study of some economic traits of seven genotypes of chickens under intensive production system. *Egypt. Poult. Sci. J.*, 20: 1031-1046.
- Pirchener, F. (1979). *Population genetik in der tierzucht*. Poult. Parey, Hamburg and Berlin.
- SAS Institute (1994). *SAS users guide: Statistics, Ver., 6.04, Fourth Edition* SAS Institute Inc., Cary NC.USA.
- Sherif, B. T. (1991). Improvement of some economical traits in chickens. Ph. D. Thesis, Minufiya Univ., Egypt.
- Sinha, S. K. and R. Khanna, (1975). Physiological, biochemical and genetic bases of heterosis. *Advan. Agron.* 27: 123-174.
- Steel , R. G. E. and G. H. Torrie (1960). *Principles and procedures of statistics*. McGraw-Hill Book Company, Inc., New York.
- Wang, A. G. and F. pirchner (1991). Heterosis of feed efficiency and reproductive traits in reciprocal crosses of laying hens. *A. B. A.*, 60: 1284.
- Zatter, O. M. M. (1994). Genetic studies in poultry, effect of crossbreeding between new local strains of chicken on some productive traits. M. Sc. Thesis, Fac. of Agric. Alex., Univ., Egypt.

العائد الوراثي و قوة الهجين لبعض صفات إنتاج البيض في خطين من دجاج النورفا

أحمد عبد الوهاب عنب ، محمد السيد سلطان ، أسامة أحمد الوشاحي ،

فاروق حسن عبده

قسم إنتاج الدواجن - كلية الزراعة - جامعة المنوفية

الملخص العربي

الهدف الأساسي من هذه التجربة هو تكوين خطين متخصصين في سلالة النورفا (خط لإنتاج البيض و آخر لوزن الجسم) باستخدام طريقة انتخاب مستويات الاستبعاد المستقلة خلال ثلاث أجيال من الانتخاب و خلط هذه الخطوط للحصول علي قوة الهجين لصفات إنتاج البيض . كان من الواضح بصفة عامة أن البدائي في الخط المنتخب لإنتاج البيض (EN) مبكرة في النضج الجنسي وتتفوق في إنتاج البيض عن تلك التي في الخط المنتخب لوزن الجسم (BW) في حين أن الدجاجات في الخط المنتخب لوزن الجسم (BW) كانت أعلى في وزن الجسم و انتجت بيض أثقل عن تلك التي في الخط المنتخب لإنتاج البيض (EN). العائد الوراثي الحقيقي لصفات عدد البيض خلال ثلاث أجيال من الانتخاب في الخط المنتخب لإنتاج البيض في دجاج النورفا يتراوح بين ١٠.٢ إلى ٢٦ بيضة بينما العائد الوراثي المتوقع لنفس الصفات في نفس الخط تراوحت بين ١.١ إلى ٢.٥ بيضة. العائد الوراثي الحقيقي لصفة العمر عند النضج الجنسي خلال ثلاث أجيال من الانتخاب في الخط المنتخب لإنتاج البيض في دجاج النورفا يتراوح بين ٧.٣ إلى ١٦.٧ يوم بينما العائد الوراثي المتوقع لنفس الصفة في نفس الخط تراوحت بين ٠.٦٦ إلى ١.٨ يوم. أوضحت النتائج في هذه الدراسة تفوق كل هجن الجيل الأول F1 لصفات إنتاج البيض علي متوسط الآباء. أوضحت الدراسة في أن كل هجن الجيل الأول F1 لصفات إنتاج البيض ظهرت به قوة هجين موجبة فيما عدا صفة العمر عند النضج الجنسي كل هجن الجيل الأول ظهرت به قوة هجين سالبة. في الهجين (EN x BW). كانت قوة الهجين لصفات إنتاج البيض (العمر عند النضج الجنسي ، وزن الجسم عند النضج الجنسي، وزن

Genetic gains and heterosis for some egg production traits.....

الجسم عن تمام النضج ، وزن البيض عند النضج الجنسي ، وزن البيض عند تمام النضج ، عدد البيض عند ٩٠ يوم من الانتاج ، عدد البيض عند عمر ٤٢ أسبوع ، عدد البيض عند عمر ٥٢ أسبوع) - ٥.١ ، ٢.٤ ، ٦.٥ ، ٦.١ ، ٣.١ ، ١٤.٧ ، ١٣.٤ ، ٩.٢ % في حين كانت قوة الهجين لنفس الصفات للهجين (BW x EN) - ٢.٥ ، ٢.٣ ، ٣.٩ ، ٥.١ ، ٢.٩ ، ١٣.٤ ، ٦.٤ ، ٣.٩ % علي التوالي.

ويمكن أن نستخلص من نتائج هذا البحث أن خلط خطين من دجاج النورفا (أحدهما محسن لصفة عدد البيض والآخر لوزن الجسم) تعتبر وسيلة فعالة لتحسين بعض صفات إنتاج البيض لدجاج النورفا.

Table (1): Mean \pm S.d. for some egg production traits in three lines of Norfa chickens (EN, BW and control) in generations 1, 2, 3 and 4.

Generation	Traits		No.	ASM	BW _{SM}	BW _M	EW _{SM}	EW _M	EN _{90d}	EN _{42wk}	EN _{52wk}
	Lines										
Generation 1	(EN)		201	147.8 \pm 6.9a	1118.8 \pm 76.4 a	1195.1 \pm 77.9 a	36.5 \pm 2.2 a	48.8 \pm 2.1 a	60.1 \pm 4.7 a	82.9 \pm 6.7 a	120.5 \pm 7.6 a
	(BW)		195	162.3 \pm 7.3 b	1369.4 \pm 82.8 b	1518.2 \pm 95.3 b	39.6 \pm 2.4 b	51.9 \pm 2.4 b	52.2 \pm 4.6 b	73.9 \pm 5.8 b	112.1 \pm 6.7 b
	cont.		114	155.1 \pm 8.1 c	1095.6 \pm 119.2c	1221.5 \pm 192.1a	35.4 \pm 4.3 c	47.2 \pm 4.5 c	49.9 \pm 9.1 c	72.2 \pm 9.6 c	107.6 \pm 15.1 c
	Mean		510	155.1 \pm 9.4	1194.6 \pm 155.9	1311.6 \pm 197.1	37.2 \pm 3.6	49.3 \pm 3.7	54.1 \pm 7.8	76.3 \pm 8.9	113.4 \pm 11.8
Generation 2	(EN)		198	143.5 \pm 3.8 a	1067.5 \pm 51.5 a	1137.1 \pm 58.5 a	35.7 \pm 1.5 a	48.3 \pm 1.2 a	65.3 \pm 3.4 a	90.1 \pm 3.2 a	128.9 \pm 3.5 a
	(BW)		198	169.2 \pm 4.5 b	1457.1 \pm 59.2 b	1621.7 \pm 73.7 b	42.4 \pm 1.7 b	54.6 \pm 1.7 b	50.3 \pm 3.3 b	71.5 \pm 4.1 b	109.4 \pm 4.3 b
	cont.		105	157.1 \pm 7.9 c	1110.3 \pm 120.4 c	1230.1 \pm 194.2 c	36.5 \pm 5.1 a	48.1 \pm 5.6 c	48.7 \pm 9.5 b	71.6 \pm 10.1 b	106.7 \pm 16.2 b
	Mean		501	156.6 \pm 11.9	1211.6 \pm 195.7	1329.6 \pm 244.9	38.2 \pm 4.3	50.3 \pm 2.4	54.8 \pm 9.2	77.7 \pm 10.6	115.1 \pm 13.4
Generation 3 parental lines	(EN)		192	136.9 \pm 1.8 a	1030.4 \pm 28.9 a	1081.8 \pm 34.8 a	35.1 \pm 0.9 a	47.6 \pm 0.7 a	70.4 \pm 1.4 a	96.6 \pm 1.7 a	135.1 \pm 2.1 a
	(BW)		189	173.1 \pm 1.6 b	1496.7 \pm 27.8 b	1724.5 \pm 29.4 b	44.2 \pm 0.8 b	56.2 \pm 0.9 b	49.4 \pm 1.8 b	70.1 \pm 1.9 b	107.2 \pm 2.2 b
	cont.		102	153.6 \pm 8.8 c	1080.7 \pm 117.6 c	1195.3 \pm 190.2 c	34.9 \pm 4.9 a	46.1 \pm 5.8 a	50.8 \pm 10.2 b	73.9 \pm 11.9 c	109.1 \pm 15.8 b
	Mean		483	154.5 \pm 15.5	1202.6 \pm 218.7	1333.9 \pm 298.8	38.1 \pm 5.1	50.1 \pm 4.9	56.9 \pm 11.2	80.2 \pm 13.4	117.1 \pm 15.1
Generation 4 cross lines	(EN \times BW)		183	151.2 \pm 7.1 a	1292.3 \pm 64.3 a	1457.6 \pm 89.1 a	41.7 \pm 1.4 a	53.4 \pm 1.6 a	67.9 \pm 6.3 a	88.8 \pm 6.5 a	125.9 \pm 7.3 a
	(BW \times EN)		179	147.2 \pm 4.3 b	1294.3 \pm 52.5 a	1494.5 \pm 80.5 b	42.1 \pm 2.5 a	53.5 \pm 3.1 a	68.7 \pm 5.1 a	94.6 \pm 4.1 b	132.3 \pm 4.2 b
	cont.		93	156.9 \pm 9.5 c	1100.7 \pm 123.4 b	1230.6 \pm 194.3 c	36.1 \pm 4.8 b	47.9 \pm 5.3 b	49.2 \pm 10.6 b	72.5 \pm 9.9 c	108.1 \pm 15.9 c
	Mean		455	151.8 \pm 7.4	1229.1 \pm 125.3	1394.2 \pm 178.1	39.9 \pm 4.3	51.6 \pm 4.4	61.9 \pm 10.9	85.3 \pm 11.9	122.1 \pm 14.9

Mean with different letters in each trait are significantly differ at 5% level.

(EN) = egg number line

(BW) = body weight line

ASM = age at sexual maturity
maturity

BW_{SM} = body weight at sexual maturity

BW_M = body weight at

EW_{SM} = egg weight at sexual maturity

EW_M = egg weight at maturity

EN_{90d} = egg number in the first 90 days after sexual maturity.
age

EN_{42wk} = egg number at 42 weeks of

EN_{52wk} = egg number at 52 weeks of age

Table (2): Analysis of variance for some egg production traits in three lines of Norfa chickens (EN, BW and control) in generations 1, 2, 3 and 4.

source of variation	d.f.	Mean of squares							
		ASM	BW _{SM}	BW _M	EW _{SM}	EW _M	EN _{90d}	EN _{42wk}	EN _{52wk}
Bet. lines	2	58966.59**	20775814.6**	32864532.3**	8588.43**	8595.34**	48029.12**	57185.93**	73655.17**
Bet. Gen.	3	2474.56**	99674.75**	643317.70**	640.35**	388.54**	6993.45**	8061.24**	7325.87**
inter Lin x Gen.	6	13930.91**	1897432.16**	3170936.22**	832.84**	725.07**	5745.74**	10786.84**	11671.24**
error	1937	38.57	7017.91	15672.99	8.24	9.02	36.75	45.04	93.33

** Significant at 1% level

(EN) = egg number line

(BW) = body weight line

ASM = age at sexual maturity

BW_{SM} = body weight at sexual maturity

BW_M = body weight at maturity

EW_{SM} = egg weight at sexual maturity

EW_M = egg weight at maturity

EN_{90d} = egg number in the first 90 days after sexual maturity.

EN_{42wk} = egg number at 42 weeks of age

EN_{52wk} = egg number at 52 weeks of age

Genetic gains and heterosis for some egg production traits.....

Table (3): Realized and expected genetic gains for some egg production traits during three generations of selection in two lines of Norfa chickens (EN and BW).

Traits Gene.	Genetic changes	ASM			BW _{SM}			BW _M			EW _M		
		(EN)	(BW)	Contr ol	(EN)	(BW)	Contr ol	(EN)	(BW)	Contr ol	(EN)	(BW)	Contr ol
G1	x	147.8	162.3	155.1	1118.8	1369.4	1095.6	1195.1	1518.2	1221.5	36.5	39.6	35.4
	Realized absolute	-7.3	7.2		23.2	273.8		-26.4	296.7		1.1	4.2	
	expected	-1.8	1.4		13.6	17.4		-4.6	12.4		0.91	1.2	
G2	x	143.5	169.2	157.1	1067.5	1457.1	1110.3	1137.1	1621.7	1230.1	35.7	42.4	36.5
	Realized absolute	-13.6	12.1		-42.8	346.8		-93.0	391.6		-0.8	5.9	
	expected	-1.2	0.99		-11.9	14.8		-6.4	12.3		-0.64	0.74	
G3	x	136.9	173.1	153.6	1030.4	1496.7	1080.7	1081.8	1724.5	1195.3	35.1	44.2	34.9
	Realized absolute	-16.7	19.5		-50.3	416.0		-113.5	529.2		0.2	9.3	
	expected	-0.66	0.47		-8.7	8.6		-5.9	6.2		0.31	0.35	

(EN) = egg number line

(BW) = body weight line

ASM = age at sexual maturity

BW_{SM} = body weight at sexual maturity

BW_M = body weight at maturity

EW_{SM} = egg weight at sexual maturity

EW_M = egg weight at maturity

Genetic gains and heterosis for some egg production traits.....

EN_{90d} = egg number in the first 90 days after sexual maturity.
age

EN_{42wk} = egg number at 42 weeks of

EN_{52wk} = egg number at 52 weeks of age

Table (3): Cont.

Traits Gene	Genetic changes	EW _M			EN _{90d}			EN _{42wk}			EN _{52wk}		
		(EN)	(BW)	Contr ol	(EN)	(BW)	Contr ol	(EN)	(BW)	Contr ol	(EN)	(BW)	Contr ol
G1	x	48.8	51.9	47.2	60.1	52.2	49.9	82.5	73.9	72.2	120.5	112.1	107.6
	Realized absolute	1.6	4.7		10.2	2.3		10.3	1.7		12.9	4.5	
	expected	0.81	1.9		2.5	2.1		2.0	1.2		1.9	1.3	
G2	x	48.3	54.6	48.1	65.3	50.3	48.7	90.1	71.5	71.6	128.9	109.4	106.7
	Realized absolute	0.2	6.5		16.6	1.6		18.5	-0.1		22.2	2.7	
	expected	0.41	0.69		2.4	2.0		2.0	-2.2		1.2	1.1	
G3	x	47.6	56.2	46.1	70.4	49.4	50.8	96.9	70.1	73.9	135.1	107.2	109.1
	Realized absolute	1.5	10.1		19.6	-1.4		23.0	-3.8		26.0	-1.9	
	expected	0.23	0.38		1.2	-1.1		1.1	-1.2		1.2	-1.2	

(EN) = egg number line

(BW) = body weight line

ASM = age at sexual maturity

BW_{SM} = body weight at sexual maturity

BW_M = body weight at maturity

EW_{SM} = egg weight at sexual maturity

EW_M = egg weight at maturity

EN_{90d} = egg number in the first 90 days after sexual maturity.

EN_{42wk} = egg number at 42 weeks of

age

A.A.Enab, M.E.Soltan, O. A. El-Weshahy and F.H. Abdou

EN_{52wk} = egg number at 52 weeks of age

Table (4): Mean performance of the parental and hybrids, heterosis from the mid-parents (MP) for egg production traits of Norfa chickens in generation four.

Cross lines Traits	(EN × BW)					(BW × EN)				
	P ₁	P ₂	MP	F ₁	ADH %	P ₁	P ₂	MP	F ₁	ADH %
ASM	136.9	173.3	155.1	151.2	-2.5	173.3	136.9	155.1	147.2	-5.1
BW _{SM}	1030.4	1496.7	1263.6	1292.3	2.3	1496.7	1030.4	1263.6	1294.3	2.4
BW _M	1081.8	1724.5	1403.2	1457.6	3.9	1724.5	1081.8	1403.2	1494.5	6.5
EW _{SM}	35.1	44.2	39.7	41.7	5.1	44.2	35.1	39.7	42.1	6.1
EW _M	47.6	56.2	51.9	53.4	2.9	56.2	47.6	51.9	53.5	3.1
EN _{90d}	70.4	49.4	59.9	67.9	13.4	49.4	70.4	59.9	68.7	14.7
EN _{42wk}	96.6	70.1	83.4	88.8	6.5	70.1	96.6	83.4	94.6	13.4
EN _{52wk}	135.1	107.2	121.2	125.9	3.9	107.2	135.1	121.2	132.3	9.2

(EN) = egg number line

P₁ = the mean of the first parent

ASM = age at sexual maturity

(BW) = body weight line

P₂ = the mean of second parent

BW_{SM} = body weight at sexual maturity

BW_M = body weight at

EW_{SM} = egg weight at sexual maturity EW_M = egg weight at maturity

EN_{90d} = egg number in the first 90 days after sexual maturity.

age

EN_{52wk} = egg number at 52 weeks of age

EN_{42wk} = egg number at 42 weeks of

