

ENHANCED FABA BEAN GROWTH BY COMBINED INOCULATION WITH *Rhizobium* STRAINS AND *Pseudomonas fluorescens* PF-23932 STRAIN AS A PLANT GROWTH PROMOTING RHIZOBACTERIA

El Sayed, A. I.; A. S. El-Sanousy and M.A. Nassef
Microbiology Department, Soil, Water and Environmental
Institute, Agriculture Research Center, Giza, Egypt.
E-mail: Dr.ahmedie@yahoo.com



ABSTRACT

Response of faba bean (*Vicia faba* L.) to co-inoculation with *Rhizobium leguminosarum* bv. *viciae* strains as a plant growth promoting and *Pseudomonas fluorescens* with half recommended doses of NPK was studied. *Rhizobium* strain RL-12612 showed significantly increased in Indole acetic acid (IAA) produced in complete and minimal medium supplemented with tryptophan in relation to bacterial strains. Whereas, PF-23932 showed significantly increased in IAA produced in minimal medium supplemented with tryptophan. A field experiment was conducted at Tag El-Ezz Research Station, Dakahlia Governorate, Egypt during two seasons 2014 to 2015. Co-inoculation with *rhizobium* strains and *Pseudomonas fluorescens* PF-23932 improved nodulation parameters during two seasons above the uninoculated plants. All treatments showed significant increase in root dry weight at 60 days plant-old above uninoculated plants during two seasons. Whereas, all treatments appeared significant increase in plant height in 2014 season. The result revealed high significant increase in the number of pods per plant in both seasons when the plants inoculated with RL-12612 strain and co-inoculated *rhizobium* strains with *Pseudomonas fluorescens* PF-23932 above uninoculated plants at 150 days-old among two seasons. RL-207 and co-inoculation with *Pseudomonas fluorescens* PF-23932 play an important role to improve chlorophyll b at 60 days plant-old among first seasons. All treatments appeared significant increase in NPK content at 60 days plant-old above uninoculated plants among two seasons except potassium content in second season.

Keywords:- Combined inoculation, IAA production, Nodulation efficiency, *Pseudomonas*, *Rhizobium*, *Vicia faba*.

INTRODUCTION

Legumes are used for a wide variety of purposes mainly as human food and animal feed. In addition to being rich in protein, legumes are also high in bone building minerals and vitamins essential for good health (FAO, 1984). Faba bean (*Vicia faba* L.) is one of the most important legume crops in Egypt. It is grown mostly to fulfill food and feed requirements for human and animal consumption. Seeds of faba bean are rich in protein (26:28%) and some other compounds. An increased production and consumption of legumes is highly desirable taking into account the high nutritional value and the beneficial health effects (Morsy *et al.* 2005).

Nitrogen (N) is the most significant yield-limiting element in many agricultural production systems (Reeve *et al.*, 2010). Nitrogen is an essential

constituent of proteins, nucleic acids, some carbohydrates, lipids, and many metabolic intermediates involved in synthesis and transfer of energy molecules (Davis, 1980). Chemical fertilizers are the most expensive energy element (Brkić *et al.* 2004). Chemical fertilizers are generally used to supply essential nutrients to the soil–plant system throughout the world. However, the prices, availability, and the environmental concerns of chemical fertilizers especially the N fertilizers are real issues of today's agriculture. Therefore, there is an urgent need to find alternative strategies that can ensure competitive crop yields, provide environmental safety, and protection while maintain long term ecological balance in agro-ecosystem (Majeed *et al.*, 2015).

Biological nitrogen fixation, one of the most important soil processes, is developed by means of microorganisms. Use of microbial inoculants or plant growth- promoting rhizobacteria (PGPR) for the enhancement of sustainable agricultural production is becoming a more widely accepted practice in intensive agriculture in many parts of the world to avoid environmental pollution resulted from chemical fertilizers (Kumar *et al.*, 2014).

A special role in biological nitrogen fixation is attributed to symbiosis between nodule bacteria of the genus *Rhizobium*, *Bradyrhizobium* and legumes. Rhizobia are the most studied PGPR for their potential to fix N₂ in the leguminous plants (Gage, 2004). Burns and Hardy (1975) reported that *Rhizobium* and *Bradyrhizobium* fixed about 175 × 10⁶ tons of nitrogen per year. It is many times higher amount of nitrogen in comparison with the world annual production of nitrogen fertilizers being 30 × 10⁶ tons (Postgate, 1982). The N₂-fixing capability of rhizobia varies greatly up to 450 Kg N/ha among host plants species and bacterial strains (Stephens and Rask, 2000).

Plant growth promoting rhizobacteria (PGPR) are a heterogeneous group of bacteria that can be found in the rhizosphere, at root surfaces and in association with roots, which can improve the extent or quality of plant growth directly and/or indirectly. In last few decades a large array of bacteria including species of *Pseudomonas*, *Azospirillum*, *Azotobacter*, *Klebsiella*, *Enterobacter*, *Alcaligenes*, *Arthrobacter*, *Burkholderia*, *Bacillus* and *Serratia* have reported to enhance plant growth (Glick, 1995). *Pseudomonas fluorescens* is adapted to survival in soil and colonization of plant roots. This applies also to the particular case of biocontrol agents from this species. The microbial inoculants that are used in agriculture include biofertilizers, biocontrol agents and plant growth promoting rhizobacteria (Kiely *et al.*, 2006). PGPR-mediated mechanisms to enhancement plant growth and yields in many crops. They are produce ACC deaminase to reduce the level of ethylene in the roots of the developing plants and thereby increasing the root length and growth. They are produced hormones like indole acetic acid (IAA), gibberellic acid and cytokinins; asymbiotic nitrogen fixation; solubilization of mineral phosphates and mineralization of other nutrients and control of phytopathogenic microorganisms (Cassan *et al.*, 2009).

Simultaneous inoculation of legumes with *Rhizobium* and free-living rhizobacteria belonging to the genera *Pseudomonas* has been reported to give positive effects on nodulation for several legumes (Oliveira *et al.* 1997).

Therefore, this study was devoted to investigating the synergistic interaction of *Rhizobium* and *Pseudomonas fluorescens* for improving faba bean growth and yield.

MATERIALS AND METHODS

Materials

Bacterial strains and growth conditions:

Bacterial strains used in this study are listed in Table 1, which including their references, as well as, their designations.

Table1. Bacterial strains used in this study.

Strains	Source or Reference	Designation
<i>Rhizobium leguminosarum</i> (12612)	IAM culture collection, Univ. of Tokyo, Japan.	RL- 12612
<i>Rhizobium leguminosarum</i> (3841)	Kindly provided by Prof J P W Young, Department of Biology, University of Yourk , UK.	RL-3841
<i>Rhizobium leguminosarum</i> (ARC 207)	Agric. Res. Center, Dept. of Microbiology, Giza, Egypt.	RL -207
<i>Pseudomonas fluorescences</i> (NRRL B-23932)	National Center for Agriculture Utilization Research, USA.	PF-23932

Media: *Rhizobium* strains were grown at 28 C in yeast extract-mannitol medium (YEM) according to Vincent (1970). It was supplemented with 0.1 mgml⁻¹ L-tryptophan for IAA assay.

Yeast extract-mannitol-congo red agar (YMCRA): This medium was used to checked rhizobium strains according to Pattison and Skinner (1974).

Kings Medium: This medium was used for the maintenance of *Pseudomonas* strains according to Merck (1994). It was supplemented with 0.5g tryptophan for determining IAA in *Pseudomonas* strains according to Ahmad *et al.* (2005).

Minimal Medium: This medium supplemented with 0.5g tryptophan to be used for IAA assay according to Balassa (1963).

M9 minimal medium: This medium used as a minimal medium for IAA determination in *Pseudomonas* strains according to Leveau and Lindow (2005).

Soil analysis: Soil samples representing the studied location (Tag El-Ezz Research Station, Dakahlia Governorate, Egypt) were analyzed before cultivation process for some physico-chemical properties (Table 2) according to Page *et al.* (1982) and Klute (1986).

Table 2. Chemical and physical characteristics of experimental soil used in this study.

Soil characteristics	Value	
	2013-2014	2014-2015

Mechanical analysis (%)		
Course sand %	3.77	3.52
Fine sand %	30.65	31.32
Silt %	30.42	40.23
Clay %	30.16	25.11
Texture class	Clay loamy	Clay loamy
Physical and chemical analysis		
pH	8.9	7.6
Organic matter %	1.45	1.32
E.C dsm ⁻¹ (1: 5) extract	1.78	1.65
Available macro and micro-nutrients		
N mg kg ⁻¹	21	24
P mg kg ⁻¹	7.92	7.21
K mg kg ⁻¹	312.5	215.21
Ca meq/100g	2	1.77
Mg meq/100g	1	0.965
Na meq/100g	5.7	4.3
K meq/100g	0.2	0.12
SO ₄ ²⁻ meq/100g	1.9	1.85

Plants : Faba bean (*Vicia faba L.*) seeds from Giza 843 were used. Seeds were obtained from Agronomy Res. Institute, Agric. Res. Centre, Ministry of Agric., Giza.

Methods

Preparation of inoculum

Yeast extract mannitol (YEB) was prepared in conical flasks and autoclaved at 121°C. This broth was then inoculated with *Rhizobium* strains and incubated at 28° C for three days. However, *Pseudomonas* strain was inoculated in conical flasks containing King's B medium according to Merck (1994). The flasks shaking at 150 rpm for 48 hrs at room temperature (25 ± 2°C). Inoculants were applied in soil with 2 mL pure culture/pot (1 × 10¹² cfu mL⁻¹).

IAA-detection with the Salkowski Colorimetric Technique

Production of IAA in the supernatant was assayed using the PC method, as described by Pilet and Chollet (1970). This method was shown to be most sensitive and most specific (Glickmann and Dessaux, 1995).

Field experiments

A field experiment was designed and conducted at Tag El-Ezz Research Station, Dakahlia Governorate, Egypt, during the winter seasons of 2013-2014 to 2014-2015. This experiment was conducted in a randomized complete block design with three replicates. Ten treatments were used as follows:

- 1- Uninoculated plants (control)
- 2- 100%N+100%P (recommended doses).
- 3- 50% N+100% p (recommended doses).
- 4- RL- 12612+ 50% N +100 %P.
- 5- RL-3841+ 50% N +100 %P.
- 6- RL- 207+ 50% N +100 %P.
- 7- PF-23932 + 50% N +100 %P.

8- RL- 12612 + PF-23932 + 50% N+100 %P.

9- RL-3841+ PF-23932 + 50%N+100%P.

10- RL-207 + PF-23932 + 50% N +100%P.

Nitrogen and phosphorus fertilizers were applied to the control treatment at the rate of 15 kg/fed ammonium sulphate (20.5%N) and 25150 kg/fed superphosphate (15.5% P₂O₅), respectively. These additions were as recommended by Egyptian Ministry of Agriculture.

Plant sampling

Plants were harvested in two stages 60-days-plant old and harvesting time.

Nodulation

Nodules developed on the plants were counted after 60-days-plant old of inoculation. Three plants from each replicate were removed and washed carefully with tap water. Nodule number were counted, dried and weighted according to Novak *et al.* (2004). Average weight of nodule (AWON) was estimated according to Pereira *et al.* (1989).

Vegetative Traits

Different parts of plants (shoots and roots) at 60-days-plant-old were oven dried at 70°C until reached to a constant mass and then turned immediately to weight. Plant height was measured when the plants became to blooming at harvest time by centimeters from the first leaf to the apex. Leaf area/plant was determined using leaf fresh weight method according to Beadle (1993). At harvesting time total NPK content, number of pods/plants and weight of one hundred seed were determined.

Plant mineral content analysis

Oven – dried samples of roots, shoots and seeds were powdered for estimation of total N, P and K⁺. Estimation of total N was done by kjeldahl's method, while the total P content was measured by vanado – molybdate yellow color method (Papakosta and Gagianas, 1991). Total K⁺ were measured by flame photometric method (Ryan *et al.*, 2001).

Photosynthetic Pigments:

Chlorophyll concentration (chl. a, b, total and carotene per g tissue) in plant leaves was extracted in 80% methanol. The pigments were determined spectrophotometrically after stored the extracted solution for twenty four hours in a refrigerator and calculated according to the Lichtenthaler and Wellburn (1983).

Total protein assay

Protein was extracted from seeds at harvesting time by dilute alkaline hydrolysis. Then proteins in the supernatants were quantified by the Coomassie Brilliant Blue procedure for protein determination (Bradford, 1976) was used to determine protein concentration. However, Bovine serum albumin ranging in concentrations from 0 to 100 µg/ml was used as the standard from the standard curve.

Free amino acids assay

Free amino acids were determined in plant leaves at 60-days-plant-old according to the method of Lee and Takahashi (1966).

Determination of total phenolic content

Total phenolics were determined using the Folin±Ciocalteau reagent according to Singleton & Rossi (1965).

Statistical analysis:

Data were subjected to the analysis of variance according to Snedecor and Cochran (1955). Least significant difference (L.S.D.) was used to compare between means if the F-test was significant maintained for further experimentation.

RESULTS AND DISCUSSION

Indole-3-acetic acid production

Indole acetic acid (IAA) production is a major property of rhizosphere bacteria that stimulate and facilitate plant growth (Mohite, 2013). In the present investigation bacterial strains were evaluated based upon quantitative estimation of IAA in complete and minimal medium in presence of tryptophan to selected the efficient ones to be used as a plant growth promoting biofertilizers. As shown from the results presented in Fig1, RL-12612 showed significantly increased in IAA produced in complete and minimal medium in relation to other strains. Whereas, PF-23932 showed significant production of IAA produced in minimal medium. IAA production varied between 27.18 to 7.89 in complete medium and 2.3 to 7.01 in minimal medium. This result agreed with Mohite (2013), who found that five rhizobacterial isolates preferred tryptophan for production and the maximum production was found in the medium amended with 0.1% tryptophan. Garge *et al.* (2015) found that all *Rhizobium* isolated from the root nodules of berseem produced IAA after 72 h, however their production varied significantly. Williams and Signer (1990) found that alfalfa symbiont *Rhizobium meliloti* produces indole-3-acetic acid in a regulated manner when supplied with exogenous tryptophan. The strains differ in their growth and production of IAA on different carbon and nitrogen sources (Sridevi and Mallaiah 2007). Indole acetic acid (IAA) is one of the most physiologically active auxins and common product of L-tryptophan metabolism by several microorganisms including rhizobia (Mandal *et al.* 2007).




Figure1. Qualitative assessment of IAA production by *R. leguminosarum* strains and *Pseudomonas fluorescences*.

Symbiotic effectiveness

As shown from the results presented in the Table 3, co-inoculation (RL-12612+ PF-23932, RL-3841+ PF-23932 and RL-207+ PF-23932) improved nodulation parameters during two seasons above the uninoculated plants. *Rhizobium* strains (RL-3841 and RL-207) appeared significant increase in nodules number developed per plant in the second season above the uninoculated plants. Whereas, *Rhizobium* strains (RL-12612, RL-3841) appeared significant increase in nodules dry weight per plant during two seasons and nodulation index in the second season. On the other hand, *rhizobium* strain RL-207 and co-inoculation (RL-207 + PF-23932) appeared significant increase in average weigh of nodule above the uninoculated plants. These results agreed with Fatma *et al.*(2012), who found that the combination of AM fungi and *Rhizobium tropici* induced a significant increase in nodulation in common bean (*Phaseolus vulgaris* L.). Smith and Zhang (1999) reported that inoculating soybean with preinduced *Bradyrhizobium japonicum* improved soybean nodulation. Thilak *et al.* (2006) found that *Rhizobium* can also affect the growth and nitrogen fixation in pigeon pea by enhancing the occupancy of introduced *Rhizobium* in the nodules of the legume. The results indicated that nodulation index of *Rhizobium* strains with *Pseudomonas putida* was higher than that of *Rhizobium*.

Table 3. Means of nodulation parameters in faba bean plants inoculated with *Rhizobium*, *Pseudomonas*.

Treatments	Nodules developed /plant		Nodule DW† (mg/plant)		Average weight of nodule (mg)		Nodulation index	
	I	II	I	II	I	II	I	II
Uninoculated plants	2	4	0.004	0.005	0.003	0.001	0.074	0.052
Full dose of N	5	3	0.005	0.009	0.001	0.002	0.037	0.063
Half dose of N	6	3	0.067	0.017	0.011	0.005	0.567	0.147
RL-12612	18	3	0.106	0.155	0.006	0.079	0.728	1.559
RL-3841	17	9	0.136	0.163	0.008	0.020	0.907	1.429
RL-207	20	13	0.060	0.092	0.026	0.044	0.484	0.879
PF-23932	2	2	0.002	0.004	0.0001	0.0003	0.016	0.037
RL-12612+ PF-23932	24	13	0.137	0.179	0.005	0.014	1.140	1.203
RL-3841+ PF-23932	30	17	0.189	0.189	0.007	0.011	1.749	1.594
RL-207+ PF-23932	18	13	0.184	0.155	0.012	0.012	1.274	1.143
F-test	NS	*	*	*	*	NS	*	**
LSD								
0.05		3.69	0.08	0.10	0.01		0.61	0.92
0.01		5.43	0.11	0.14	0.01		0.84	1.27

NS, ** = Insignificant differences , significantce at 0.05 and 0.01 probability levels, respectively.

†= Dry weight.

I,II = Growing seasons of 2013/2014 and 2014/2015, respectively.

The data presented in Table 4 showed that plants inoculated with RL-3841+PF-23932 showed significant increase in root dry weight above plants fertilized with recommended dose. However, all inoculum were expressed significant increase in shoot dry weight above uninoculated plants. Whereas, all inoculum appeared significant increase in plant height in the second season. This may be due to environmental condition of this year which stimulated more production of indole-3-acetic acid (IAA) leading to play a significant role in plant growth (Newcomb, 1980). These result agreed with El-Wakeil and El-Sebai (2009), who found a significant positive effect of rhizobium strains on fresh and dry weight of leaves and stems, root/ shoot ratio, pods/flowers ratio, as well as, the number and weight of nodules compared to NPK fertilizer plots. Ögutcu *et al.* (2008) found significant differences among rhizobial strains for various parameters such as nodule dry weight and shoot dry weight. Deshwal, *et al.* (2003) reported that rhizobia are known to increase nodulation and nodule dry weight in legumes along with increase in host plant growth and development. Rudresh *et al.* (2004) found that a combined inoculation of *Rhizobium* and a phosphate solubilizing *Bacillus* increased germination, nutrient uptake, plant height, number of branches, nodulation, yield and total biomass of check pea if compared to either individual or uninoculated control.

Table 4. Mean of growth performance in inoculated plants.

Treatments	Root DW ^T (g/plant)		Shoot DW [†] (g/plant)		L.A/P ^T (cm ²)		Plant height (cm)	
	I	II	I	II	I	II	I	II
Uninoculated plants	0.6	2.7	6.5	9.4	844	748	63	59
Full dose of N	3.0	2.8	14.3	13.6	3070	3333	85	81
Half dose of N	2.4	1.8	12.3	11.7	1295	2553	79	75
RL-12612	2.1	1.0	14.7	10.1	2619	2054	74	78
RL-3841	1.9	1.2	15.0	11.6	2971	2627	71	77
RL-207	2.0	1.4	11.6	11.2	2120	1690	75	70
PF-23932	1.8	0.9	14.6	10.0	4215	2694	72	75
RL-12612+ PF-23932	2.9	2.6	12.6	15.0	3909	4364	71	79
RL-3841+ PF-23932	3.4	2.7	11.3	12.3	2182	1375	77	82
RL-207+ PF-23932	3.3	2.5	15.0	13.7	3053	2782	85	91
F-test	*	*	*	*	**	*	**	**
LSD 0.05	0.37	0.53	4.8	2.13	3055.66	2209.53	12.89	10.35
0.01	0.51	0.73	6.58	2.92	4190.62	3030.22	17.67	14.19

*,** = Significant at 0.05 and 0.01 probability levels, respectively.

†= Leaf area per plant. I, II= Growing seasons of 2013/2014 and 2014/2015, respectively.

Yield parameters

Data presented in Table 5 showed the effect of inoculation on yield components (number of pods per plant and seeds dry weight) among two seasons. The results revealed a significant increase in the number of pods developed per plant above uninoculated plants in both seasons when plants inoculated with RL-12612 and co-inoculation (RL-12612+ PF-23932, RL-3841+ PF-23932 and RL-207+ PF-23932). Whereas, *Rhizobium* strains RL-3841 and RL-207 appeared significant increase in the number of pods per plant in 2013/2014 season and seeds dry weight among both seasons. This result agreed with Fatnassi *et al.* (2015), who found that *Vicia faba* inoculated with *Rhizobium leguminosarum* bv. *Viciae* appeared significant increases the number and the weight of nodules by 50%. Whereas, co-inoculation between *Rhizobium* and *Pseudomonas* positively influence the growth and seed yield per plant. Dashadi *et al.* (2011) found that co-inoculation of *Rhizobium* and *Azotobacter* increased total nitrogen content, nodulation, seed yield and biological yield under water deficit condition. However, El-Wakeil and El-Sebai (2009) found that the highest number of pods was achieved in treatment of rhizobia mixed with mycorrhiza or *pseudomonas*. Several workers have reported that seed inoculation with *Rhizobium* has significantly increased the growth and yield of legume crops (Pathak *et al.*, 2001).

Table 5. Means of yield parameters

Treatments	Pods/plant		Seeds (g/plant)	
	I	II	I	II
Uninoculated plants	12.33	11.33	18.20	25.95
Full dose of N	31.67	27.33	52.07	58.28
Half dose of N	22.67	21.00	36.60	27.68
RL-12612	22.00	19.33	43.01	40.21
RL-3841	20.00	18.00	41.23	42.20
RL-207	13.33	20.67	37.74	39.71
PF-23932	11.00	15.00	26.83	32.56
RL-12612+ PF-23932	25.00	23.67	51.91	50.92
RL-3841+ PF-23932	23.33	24.00	46.89	57.97
RL-3841+ PF-23932	32.33	25.67	58.99	57.13
F-test	*	**	*	**
LSD 0.05	8.29	6.36	8.88	8.59
0.01	11.37	8.72	12.18	11.78

*, ** = Significant at 0.05 and 0.01 probability levels, respectively.
I, II = Growing seasons of 2013/2014 and 2014/2015, respectively.

Biochemical traits

Data presented in Table 6 showed that RL-207 and co-inoculation (RL-12612+ PF-23932, RL-3841+ PF-23932 and RL-207+ PF-23932) play an important role to improve chlorophyll b at 60 days plant-old at the first season. These results agreed with Kosslak and Bohlool (1984), who found that photosynthetic capacity of the host plant may be affected by number of successful infections. Benefits to plants from PGPR have been shown to include increases in seed germination rate, root growth, yield, leaf area, chlorophyll content, nutrient uptake, protein content, hydraulic activity, tolerance to abiotic stress, shoot and root weights, biocontrol, and delayed senescence (Yang *et al.* 2009). Dakora (2003) reported that the release phytohormones produced by rhizobia into cropping systems promoted plant growth through N₂ fixation.

Table 6. Chlorophyll pigments in the leaves of faba bean plants.

Treatments	Chlorophyll content (mg/g)						Carotene	
	Chl [†] .a		Chl.b		Total Chl.		I	II
	I	II	I	II	I	II		
Uninoculated plant	1.45	1.58	0.23	0.43	1.68	2.01	1.42	1.28
Full dose of N	1.63	2.87	0.23	0.59	1.87	3.46	1.41	2.21
Half dose of N	2.12	2.40	0.23	0.43	2.35	2.83	1.90	1.57
RL-12612	1.77	1.80	0.24	0.36	2.01	2.16	1.90	1.64
RL-3841	2.98	1.80	0.45	0.42	3.43	2.23	2.32	1.41
RL-207	2.84	1.80	0.61	0.40	3.45	2.21	2.24	1.30
PF-23932	2.00	1.64	0.52	0.35	2.53	1.99	1.84	1.27
RL-12612+ PF-23932	3.06	1.49	0.75	0.37	3.81	1.86	2.41	1.15
RL-3841+ PF-23932	2.66	2.14	0.64	0.54	3.30	2.68	2.20	1.85
RL-207+ PF-23932	1.98	2.11	0.57	0.46	2.55	2.57	1.87	1.81
F-test	NS	*	**	*	NS	*	NS	NS
LSD 0.05		0.76	0.23	0.19		0.87		
0.01		1.04	0.31	0.27		1.20		

NS, *, ** = Insignificant differences, significant at 0.05 and 0.01 probability levels, respectively.

† = chlorophyll. I, II = Growing seasons of 2013/2014 and 2014/2015, respectively.

Data summarized in Table 7 revealed that all treatments appeared significant increase in NPK content at 60 days plant-old above uninoculated plants among two seasons except for potassium content in the second season. This result agreed with Kucuk and Kivanc (2008), who found significant increase in seed yield, seed weight and protein range in three bean (*Phaseolus vulgaris* L.) varieties inoculated by *Rhizobium* spp. The increase in dry-matter and nitrogen content of co-inoculated plants may be attributed to increased nodulation, higher N₂-fixation and improvement of root development under controlled or field conditions (Dashti *et al.* 1998). PGPR bacteria can increase P availability to plants through solubilizing insoluble phosphates and this may improve biological nitrogen fixation and availability of other nutrients (Gyaneshwar *et al.*, 2002).

Table 7 . Mean of NPK content of faba bean grown under field conditions.

Treatments	N		p		k	
	I	II	I	II	I	II
Uninoculated plants	1.3	1.3	0.10	0.12	1.15	1.16
Full dose of N	1.7	1.8	0.23	0.25	1.69	1.90
Half dose of N	1.8	1.8	0.24	0.21	1.85	1.76
RL-12612	1.9	1.9	0.21	0.22	1.60	1.69
RL-3841	2.2	1.9	0.24	0.23	1.65	1.77
RL-207	1.9	2.1	0.30	0.27	1.89	1.73
PF-23932	1.6	1.5	0.27	0.24	1.62	1.79
RL-12612+ PF-23932	2.3	2.0	0.31	0.31	1.74	1.75
RL-3841+ PF-23932	1.7	2.2	0.22	0.26	1.83	1.76
RL-207+ PF-23932	2.9	2.6	0.34	0.29	1.73	1.85
F-test	*	*	*	*	*	NS
LSD 0.05	0.36	0.19	0.06	0.04	0.26	

NS,* = Insignificant differences and significantce at 0.05 probability levels, respectively. I, II = Growing seasons of 2013/2014 and 2014/2015, respectively.

Data in Table 8 showed that all treatments induced significant increase in total protein of seeds above uninoculated plants at the first season. The same effect was achieved by RL-207 and co-inoculation (RL-12612+ PF-23932, RL-3841+ PF-23932 and RL-207+ PF-23932) at the second season. Co-inoculation (RL-12612+ PF-23932 and RL-207+ PF-23932) appeared significant increase in total free amino acids in shoot above uninoculated plants among two seasons. However, bacterial strains RL-12612, RL-3841, RL-207, PF-23932 and co-inoculation (RL-3841+ PF-23932) appeared significant increase in total free amino acids in shoots above uninoculated plants among the second season. All treatments except for half recommended dose and RL-3841 appeared significant increase in shoot phenol content above uninoculated plants at the first season. These results agreed with Yadegari and Asadi Rahmani (2010), who found that co-inoculation strains with *P. fluorescens* gave the highest seed yield, number of pods per plant, weight of 100 seed, seed protein yield, number of seed per pod, and seed protein yield of bean (*Phaseolus vulgaris*). However, Solaiman

and Rabbani (2005) found that the performance of *Rhizobium* inoculant was superior in relation to uninoculated control concerning protein content in green and mature seeds of pea.

Table 8. Protein chemical traits in faba bean plants inoculated with rhizobia and *Pseudomonas*.

Treatments	Protein % in seeds		Total free amino acids ($\mu\text{g/g}$)		Total phenol ($\text{mg } 100 \text{ g}^{-1}$ fresh weight)	
	I	II	II	I	I	II
Uninoculated plants	16.9	17.0	11.41	8.40	122.3	151.5
Full dose of N	23.6	23.6	43.09	19.29	285.3	303.9
Half dose of N	20.2	20.1	10.63	11.10	176.4	185.51
RL-12612	21.1	19.6	15.62	16.49	254.3	211.0
RL-3841	21.0	21.2	14.28	14.39	208.2	253.0
RL-207	20.0	19.9	23.94	18.17	288.2	231.8
PF-23932	19.7	18.4	16.15	21.02	244.8	212.3
RL-12612+ PF-23932	22.6	22.5	52.06	32.15	286.9	271.2
RL-3841+ PF-23932	22.4	24	23.83	24.71	264.6	294.3
RL-3841+ PF-23932	24.9	23.5	41.96	28.54	257.3	314.8
F-test	*	*	*	**	**	*
LSD						
0.05	2.72	3.68	15.44	5.66	96.82	92.96
0.01	3.73	5.05	21.17	7.76	132.78	127.49

*, ** = Significant at 0.05 and 0.01 probability levels, respectively.

I, II = Growing seasons of 2013/2014 and 2014/2015, respectively.

In conclusion, the present study demonstrated a positive effect of co-inoculation between *Rhizobium* with *Pseudomonas* on symbiotic effectiveness and chlorophyll contents in faba bean plants, which improving the yield and yield components.

REFERENCES

- Ahmad, F., I. Ahmad and M.S. Khan, 2005. Indole Acetic Acid Production by the indigenous isolates of *Azotobacter* and *Fluorescent Pseudomonas* in the presence and absence of tryptophan. *Turk. J. Biol.*, 29: 29-34.
- Balassa, G., 1963. Genetic transformation of *Rhizobium*: A review of the work of R. Balassa. *Bacteriol Rev.*, 27: 228-241.
- Beadle, C.L., 1993. Growth analysis. In: Hall, D.O., et al., (Eds.), *Photosynthesis and Production in a Changing Environment: A Field and Laboratory Manual*. Chapman & Hall, London, pp: 36–46.
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248–254.
- Brkić, S. Milaković, Z. Kristek, A. Antunović, M. 2004. Pea yield and its quality depending on inoculation, nitrogen and molybdenum fertilization. *Plant Soil Environ.*, 50, (1): 39–45.
- Burns R.C. and Hardy R.W.F. 1975. *Nitrogen fixation in bacteria and higher plants*. Springer-Verlag, New York.

- Cassan F, Maiale S, Masciarelli O, Vidalc A, Lunaa V, Ruiz O .2009. Cadaverine production by *Azospirillum brasilense* and its possible role in plant growth promotion and osmotic stress mitigation. *Europ J Soil Biol* 45:12–19.
- Dakora, F.D. 2003. Defining new roles for plant and rhizobial molecules in sole and mixed plant cultures involving symbiotic legumes. *New Phytol.* 158: 39 – 49.
- Dashadi, M.; Khosravi, H. ; Moezzi, A.; Nadian, H. ; Heidari, M. and Radjabi, R. 2011. Co Inoculation of *Rhizobium* and *Azotobacter* on Growth of Faba bean under Water Deficit Conditions *American-Eurasian J. Agric. & Environ. Sci.*, 11 (3): 314-319.
- Dashti N, Zhang F, Hynes R, and Smith DL .1998. Plant growth promoting rhizobacteria accelerate nodulation and increase nitrogen fixation activity by field grown soybean [*Glycine max* (L.) Merr.] under short season conditions. *Plant Soil* 200: 205-213.
- Davis, L.C. 1980. Limiting factors in nitrogen fixation. *What's New in Plant Physiology* 11(11):41 45.
- Deshwal, V.K.; R.C. Dubey and Maheshwari, 2003. Isolation of plant growth promoting strains of *Bradyrhizobium* (*Arachis*) sp. with biocontrol potential against *Macrophomina phaseolina* causing charcoal rot of peanut. *Curr. Sci.* 84 (3): 443-44.
- El-Wakeil, N. E. and El-Sebai, T. N.2009. Role of biofertilizer on faba bean growth, yield, and its effect on bean aphid and the associated predators, *Archives of Phytopathology and Plant Protection*, 42:12, 1144-1153.
- FAO .1984. Legumes Inoculants and their Use. A Pocket Manual. Food and Agriculture Organization. Rome, Italy.
- Fatma, T.;Mustapha, T. and Jean-Jacques D.2012. Combined inoculation with *Glomus intraradices* and *Rhizobium tropici* CIAT899 increases phosphorus use efficiency for symbiotic nitrogen fixation in common bean (*Phaseolus vulgaris* L). *Saudi Journal of Biological Sciences* Vol 19, Issue 2, 157–163.
- Fatnassi, I. C.; Chiboub, M. ; Saadani, O. ; Jebara M. and Salwa H. J.2015. Phytostabilization of moderate copper contaminated soils using co-inoculation of *Vicia faba* with plant growth promoting bacteria. *J. Basic Microbiol.* 55, 303–311.
- Gage, DJ .2004. Infection and invasion of roots by symbiotic, nitrogen-fixing rhizobia during nodulation of temperate legumes. *Microbiol Mol Biol Rev* 68:280–300.
- Garg, Varsha , Kamlesh K., Rajesh, G. and Ankit Singla. 2015. Production of indole-3-acetic acid by berseem (*Trifolium alexandrinum* L.) rhizobia isolated from Haryana, India. *Agric. Sci. Digest.*, 35: (3) 229-232.
- Glick, B.R., 1995. The enhancement of plant growth by free living bacteria. *Can. J. Microbiol.* 41, 109–114.
- Glickmann, E. and Y. Dessaux, 1995. A critical examination of the specificity of the Salkowski reagent for indolic compounds produced by phytopathogenic bacteria. *Appl. Environ. Microbiol.*, 61(2): 793-796.

- Gyaneshwar, P., G. N. Kumar, L. J. Parekh and P. S. Poole. 2002. Role of soil microorganisms in improving P nutrition of plants. *Plant Soil* 245:83-93.
- Kiely PD, Haynes JM, Higgins CH, Franks A .2006. Exploiting new systems-based strategies to elucidate plant- bacterial interactions in the rhizosphere. *Microbiol. Ecol.* 51: 257–266.
- Klute, A., 1986. *Methods of Soil Analysis: part I. physical and mineralogical methods* (2nd Ed.) Amer. Soc. Agron. Monograph No. 9. Madison-Wisconsin, USA. Monograph No. 9. Madison-Wisconsin, USA.
- Kosslak, R.M. and B.B. Bohlool, 1984. Suppression of nodule development of one side of a split root system of soybean caused by prior inoculation of the other side. *Plant Physiol.*, 75:125-130.
- Küçük, C. and M. Kıvanç. 2008. Preliminary characterization of *Rhizobium* strains isolated from chickpea nodules. *African J. Biotechnol.* 7 (6), pp. 772-775.
- Kumar A., Maurya B. R., and Raghuvanshi R. 2014. Isolation and characterization of PGPR and their effect on growth, yield and nutrient content in wheat (*Triticum aestivum* L.). *Biocatal. Agric. Biotechnol.* 3 121–128.
- Lee YP and Takanashi T.1966. An improved colorimetric determination of amino acids with the use of ninhydrin. *Anal. Biochem.*, 14: 71-77. Linchtenthaler
- Leveau, J.H.J. and S.E. Lindow. 2005. Utilization of the plant hormone indole-3-acetic acid for growth by *Pseudomonas putida* strain 1290. *Appl. Environ. Microbiol.* 71(5) : p 2365-2371.
- Lichtenthaler, H.K. and A.R. Wellburn, 1983. Determination of total carotenoids and chlorophylls a and b of leaf extracts in different solvents. *Biochem. Soc. Transactions*, 11: 591-592.
- Majeed, A.; Kaleem A. A. M., Hameed, S., Asmalman and Nasir, R. 2015. Isolation and characterization of plant growth promoting rhizobacteria from wheat rhizosphere and their effect on plant growth promotion *Plant-Microbe Interaction*, a section of the journal *Frontiers in Microbiology*. 6:198.
- Mandal S.M., K.C. Mondal, S. Dey, and B.R., Pati .2007. Optimization of cultural and nutritional conditions for indole-3-acetic acid (IAA) production by a *Rhizobium* sp. isolated from root nodules of *Vigna mungo* (L.) Hepper. *Res. J. Microbiol.* 2: 239-246.
- Merck., 1994. *Microbiology manual E*, Merck Darmstadt, Bereich Labor, D-64271, Darmstadt, Germany.
- Mohite B.2013. Isolation and characterization of indole acetic acid (IAA) producing bacteria from rhizospheric soil and its effect on plant growth. *Journal of Soil Science and Plant Nutrition* , 2013, 13(3), 638-649.
- Morsy, K.M. and A.M. Tarrad, 2005. Effect of infection with *Botrytis fabae* sard. and mechanical leaf defoliation on yield loss in faba bean. *Egypt. J. Appl. Sci.*, 20 (11B): 443-445.

- Newcomb, W. 1980. Control of morphogenesis and differentiation of pea root nodules, p. 87-102. In W. E. Newton and W. H. Orme-Johnson (ed.), Nitrogen fixation, vol. 2. University Park Press, Baltimore.
- Novak, K., L. Ludmila and S. Vlandimir, 2004. Rhizobia nod gene-inducing activity in pea nodulation mutants: dissociation of nodulation and flavonoid response. *Physiologia Plantarum*, 120: 546-555.
- Ogutcu, H. O. F. Algur, E. Elkoca and F. Kantar. 2008. The determination of symbiotic effectiveness of *Rhizobium* strains isolated from wild chickpeas collected from high altitudes in Erzurum. *Turk. J. Agric.* 32:241-248.
- Oliveira, F.L., E.M. Ferreira and M.E. Pampulha, 1997. Nitrogen fixation, nodulation and yield of clover plants co-inoculated with root-colonizing bacteria. *Symbiosis*, 23: 35-42.
- Page, A.L., R.H. Miller and D.R. Keeny, 1982. "Methods of Soil Analysis" part II. Chemical and microbiological properties (2nd ed.) Am. Soc. Agron.
- Papakosta, D.K. and Gagianas, A.A .1991. Nitrogen and dry matter accumulation, remobilization and losses for Mediterranean wheat during grain filling. *Agron J* 83: 864-870.
- Pathak, K.; Kalita, MK; Barman, U.; Hazarika, BN and Saha, N.N. 2001 Response of summer green gram (*Vigna radiata*) to inoculation and nitrogen levels in Barak Valley zone of Assam *Annals of Agri. Res.*22: 123-124.
- Pattison, A.C. and F.A. Skinner, 1974. The effects of antimicrobial substances on *Rhizobium* spp. And their use in selective media. *J. Appl. Bact.*, 37: 239-250.
- Pereira, P.A.A., R.H. Burris and F.A. Bliss, 1989. N-dinitrogen fixation potential of genetically divers 15 bean (*Phaseolus vulgaris* L.). *Plant and Soil.*, 120: 171-179.
- Pilet, P.E. and R. Chollet, 1970. Sur le dosage colorimetrique de l'acide indolyacetique. *C. R. Acad. Sci. Ser. D.*, 271: 1675-167.
- Postgate J.R. 1982. The fundamentals of nitrogen fixation. Cambridge Univ. Press., Cambridge, London.
- Reeve, w., Chain, P., O'Hara, G., Ardley, J., Nandesena, K., Bräu, L., Tiwari, R., Malfatti, S., Kiss, H., Lapidus, A., Copeland, A., Nolan, M., Land, M., Hauser, L., Chang, Y., Ivanova, N., Mavromatis, K., Markowitz, V., Kyrpides, N., Gollagher, M., Yates, R., Dilworth, M., Howieson, J., 2010. Complete genome sequence of the *Medicago* microsymbiont Ensifer (*Sinorhizobium*) medicae strain WSM419. *Stand. Genomic Sci.* 2, 77–86.
- Rudresh, D.I. ; Shivaprakash, M.K. and Prasad, R.D.2004.Effect of combined application of *Rhizobium*, phosphate solubilizing bacterium and *Trichoderma* spp. on growth, nutrient uptake and yield of chickpea (*Cicer aritenium* L.). *Appl Soil Ecol.* 28(2):139-146.
- Ryan J, Estefan G, Rashid A .2001. Soil and analysis laboratory manual. ICARDA, pp: 172.

- Singleton, V. L. and Rossi, J. A. 1965. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am. J. Enol. Vitic.*, 16: 144-158.
- Smith D. L .and F. Zhang .1999. Composition for enhancing grain yield and protein yield of legumes grown under environmental conditions that inhibit or delay nodulation thereof. U.S. Patent.5,922,316.
- Snedecor, G.W. and W.G. Cochran, 1955. *Statistical Methods*, sixth edition. The Iowa state University Press, Ames, Iowa, U.S.A.
- Solaiman, A. R. M. and M. G. Rabbani. 2005. Effects of *Rhizobium* inoculant and Nitrogen Application of Pea. *Bangladesh J. Microbiol.* 21(1): 36-40.
- Sridevi M. and K. V. Mallaiah. 2007. Bioproduction of indole acetic acid by *Rhizobium* strains isolated from root nodules of green manure crop, *Sesbania sesban* (L.) Merr. *Iranian J.Bio.*, 5, (3)178-182.
- Stephens, J. H. G. and Rask, H. M. 2000. Inoculant production and formulation. *Field Crops Research*, (65): 249-258.
- Thilak, K. V. B. R., N., Ranganayaki, and C. Manoharachari. 2006. Synergistic effects of plant-growth promoting rhizobacteria and *Rhizobium* on nodulation and nitrogen fixation by pigeonpea (*Cajanus cajan*). *European Journal of Soil Science*, 57, 67–71.
- Vincent, J.M., 1970. A manual for the practical study of root-nodule bacteria, p. 145. In I.B.M. Handbook No. 15. Blackwell Scientific Publications, Oxford, England.
- Williams, M. N. V. and E. R. Signer. 1990. Metabolism of tryptophan and tryptophan Analogs by *Rhizobium meliloti*. *Plant Physiol.* 92,1009-1013.
- Yadegari, M. and H. Asadi Rahmani.2010. Evaluation of bean (*Phaseolus vulgaris*) seeds' inoculation with *Rhizobium phaseoli* and plant growth promoting Rhizobacteria (PGPR) on yield and yield components. *African Journal of Agricultural Research* Vol. 5(9), pp. 792-799.
- Yang, J.W.; Kloepper, C.M. and Ryu.2009. Rhizosphere bacteria help plants tolerate abiotic stress. *Trends Plant Sci*, 14 ,pp. 1–4.

تعزيز نمو محصول الفول البلدى بواسطة التلقيح المشترك بين سلالات الرايزوبيم و سلالة السيدوموناس فلوروسنز PF-23932 كسماد حيوى محفز لنمو النباتات أحمد إبراهيم السيد ، أحمد سامى عمر أحمد السنوسى و محمود عبد المقصود ناصف قسم الميكروبيولوجيا الزراعية-معهد بحوث الأراضى و المياه و البيئة -مركز البحوث الزراعية.

استخدمت فى هذه الدراسة اربعة سلالات بكتيرية . تم تقييم السلالات المستخدمة بالنسبة لإنتاجها من هرمون النمو اندول حمض الخليك فى بيئى النمو الكاملة و الحديه المدعومة بالحمض الأميى تريتوفان. أظهرت سلالة الرايزوبيم RI-12612 زيادة معنوية فى إنتاج أندول حامض الخليك على مستوى بيئة النمو الكاملة و الحدية مقارنة بباقي السلالات البكتيرية. بينما أظهرت سلالة PF-23932 زيادة معنوية فى إنتاج أندول حامض الخليك على مستوى بيئة النمو الحدية مقارنة بباقي السلالات البكتيرية . تم زراعة التجربة الحقلية فى المزرعة البحثية بمحطة تاج العز للبحوث الزراعية خلال موسمي ٢٠١٣/٢٠١٤، ٢٠١٤/٢٠١٥. أظهر التلقيح المشترك بين سلالات الرايزوبيم و سلالة السيدوموناس فلوروسنز PF-23932 زيادة معنوية فى الصفات الخاصة بتكوين العقد البكتيرية على جذر نبات الفول البلدى خلال موسمين النمو . كما أظهرت كل المعاملات المستخدمة زيادة معنوية فى الوزن الجاف للمجموع الجذرى مقارنة بالنباتات الغير ملقحة خلال موسمى الزراعة ، بينما كان نفس التأثير بالنسبة لصفة طول النبات خلال موسم النمو الأول. أظهرت السلالة RL-12612 ، معاملات التلقيح المشترك بين سلالات الرايزوبيم و سلالة السيدوموناس فلوروسنز PF-23932 زيادة معنوية فى عدد القرون فاقت النباتات الغير ملقحة. لعبت سلالة الرايزوبيم RL-207 و معاملات التلقيح المشترك بين سلالات الرايزوبيم و سلالة السيدوموناس فلوروسنز PF-23932 دورا هاما فى تحسين نسبة كلوروفيل b فى الاوراق عند عمر ٦٠ يوم خلال موسم النمو الأول بينما أظهرت المعاملات زيادة معنوية فى محتوى الاوراق من المعادن النيتروجين ، الفوسفور ،البوتاسيوم خلال موسمى النمو (ماعد عنصر البوتاسيوم خلال الموسم الثانى فقط) مقارنة بالنباتات الغير ملقحة . عموما اوضحت النتائج إمكانية الإعتماد على استخدام نصف الجرعات من التسميد الكيماوى للنيتروجين مع الجرعة الكاملة للفوسفور مع استخدام التلقيح المشترك لبكتريا العقد الجذرية الرايزوبيم و السيدوموناس كمخصب حيوى فعال .